



BROOKHAVEN NATIONAL LABORATORY

ASSOCIATED UNIVERSITIES, INC., UPTON, L.I., N.Y. 11973

401367

MEDICAL DEPARTMENT

TELEPHONE: (516) 345-3577

September 2, 1976

Dr. W. W. Burr, Jr.
Deputy Director
Division of Biomedical &
Environmental Research
U.S. Energy Research and
Development Administration
Washington, D.C. 20545

Dear Bill,

With regard to the plutonium problem on Bikini we would like to bring you up to date on (1) sample collections planned for Sept.-Oct. survey in the Marshalls; (2) samples still being analyzed from previous surveys, and (3) summary Tables of previous data comparing Rongelap and Utirik findings.

Samples for Sept.-Oct. 1976: Urines 1. Bikini: 5 gal. carboys on 5 individuals (carboys distributed last June); also 5 - 5 gal. pooled samples; 2. Rongelap: 5 - 5 gal pooled samples; 3. Utirik: 5 - 5 gal. pooled samples; 4. Wotje: 5 - 5 gal. pooled samples; 5. Ebeye: 5 - 5 gal. pooled samples; 6. Majuro: 5 - 5 gal. pooled samples and samples on 50 Bikini people planning to return in the near future. You will note we are collecting samples from a number of atolls not directly involved in the 1954 fallout.

Whole body gamma spectroscopy on Bikini people (for ¹³⁷Cs body burdens) originally planned for October are being delayed until April in hope that the advance group of Eniwetok people will have returned to their atoll and can also be counted during that operation.

N. Greenhouse, J. Naidu and V. Nelson plan on collecting soil and plant samples in Sept.-Oct. from Wotje, Ailuk, Utirik, Rongelap and Bikini.

Samples presently being analyzed are animal samples (pig, chickens, crabs from Rongelap and Bikini, water from Bikini cisterns, soil from Rongelap. Autoradiographs for alpha tracks of animal lung and liver tissues are being run.

The enclosed Tables have been compiled by N. Greenhouse, J. Naidu and me and include data from various sources. Many of the Tables are incomplete. We are attempting to pull together in one file all available data from personnel and environmental radiological surveys on Rongelap, Utirik and Bikini atolls. When this is accomplished we should be able to make more comprehensive comparative Tables on these atolls. We hope these data will be on hand for your planned meeting of the experts.

Sincerely,

Bob

Robert A. Conard, M.D.

RAC:gc

Enc.

CC: J. Deal
T. McCraw
R. Ray
N. Greenhouse
J. Naidu

5009761

RADIOCHEMICAL ANALYSES OF URINE (DATA IN AVERAGE pCi/liter)^{*}

Year	No. in group	Av. vol. ml	⁹⁰ Sr	¹³⁷ Cs	²³⁹⁻²⁴⁰ Pu
<u>Rongelap</u>					
1970	20	895	3.5	2700	
1971	15	534	3.7	2400	
1972	18	460	2.4	2600	
1973	11	249	6.5	4600	
1974	14	557	2.8	4500	
1975 (Mar)	14	753	4.6	2100	
1976 (Mar)	17**	679	4.6	2100	0.009 ± .002
	1	1250			0.014 ± .007
<u>Utirik</u>					
1974	11	542	1.3	1300	
<u>Bikini</u>					
1970	Pooled		1.2	1150	
	Urine G	1100	2.2		
	Urine M	930	1.9		
	HASL * control	3000	1.0	120	
	HASL control	1000	1.6		
1971	Pooled	2670	1.7	1830	
1972	Pooled	2700	4.2	0910	
1973	13	304	6.7	1300	
1974	8	165	2.3	1300	
	10	649			
1975 (Mar)	8	360	7.3	1800	
(Oct)	18	510	3.1	1300	
	Pooled	9319			0.01 ± .002
1976 (Mar)	24**	480	4.7 ± 4.6	1600 ± 800	0.009 ± .002
<u>N.Y. City</u> (1976)					
	Pooled	20,000			0.0009 ± .0004
	1	500	0.9	8.0	
	1	500	0.8	10.0	

* Analytical error terms associated with ⁹⁰Sr and ¹³⁷Cs analyses were usually less than 10 percent.

** Pooled for Pu analysis.

5609762

11205

INDIVIDUAL RADIOTOLOGICAL MONITORING DATA - RTKINT

(Urine Levels - amt/l; ^{137}Cs Body Burdens - 1974, nCi)

No.	Age	Sex	1973			1974			1975 (Mar.)			1975 (Oct.)			1976 (Mar.)			1976 (Oct.)			
			^{90}Sr pCi	^{137}Cs nCi	Yrs. Bik.	^{90}Sr pCi	^{137}Cs nCi	^{137}Cs nCi	** Bik.	^{90}Sr pCi	^{137}Cs nCi	Bik.	^{90}Sr pCi	^{137}Cs nCi	Bik.	^{90}Sr pCi	^{137}Cs nCi	Bik.	^{90}Sr pCi	^{137}Cs nCi	Yrs. Bik.
B1	46	F				<0.1	0.5	58	1.5												
B2	20	F														2.2	2.0	1			
B3			5.5	2.6	?			72													
B4	62	M						78	2	4.5	2.1	3									
E			18.9	1.3																	
B6			2.0	0.4																	
B7	24	M						95	1												
B8	31	F						108	3												
B9	16	F														8.2	2.9	1			
10	26	M								4.7	1.9	1.5									
11	32	M				3.2	1.7	156	6												
12	51	M						290	0.5	12.0	2.1	1.5				2.9	0.9	2			
13	32	M	2.2	0.6	1			81	2							2.1	0.8	3.5			
14	24	M				<0.2	1.5		2							5.7	2.1	3			
15	57	M						51	2										2.3	0.3	4
1	25	F																			
17	56	F														4.8	2.5	1			
18	47	F						116	2												
19	M					2.4	0.8		3												
20	25	F						73	3										2.4	1.2	5
21			5.7	1.1																	
22	32	M						103	4												
23	24	M								6.6	1.5	1.5									
24	41	M	7.8	2.0	1			122	2	18.0	3.0	3				2.4	0.4	2.5	1.9	2.0	3
25	44	M														6.8		4			

5009163

INDIVIDUAL RADIOLOGICAL MONITORING DATA - BIKINI

(Urine Levels - amt/l; ^{137}Cs Body Burdens - 1974, nCi)

No.	Age	Sex	1973				1974				1975 (Mar.)				1975 (Oct.)				1976 (Mar.)				1976 (Oct.)						
			^{90}Sr pCi	^{137}Cs nCi	Yrs. Bik.		^{90}Sr pCi	^{137}Cs nCi	^{137}Cs nCi	** Yrs. Bik.	^{90}Sr pCi	^{137}Cs nCi	Yrs. Bik.																
B26	32	F									7.8	3.0	2		1.6		1.2	2.5		4.0		1.4	3						
B27	18	M									76	0.5																	
B28	15	F									77	1																	
B29	56	M									43	0.5																	
B30	33	F									252	1																	
B31	46	M																											
B32	59	M																											
B33	44	F									158	1																	
B34	49	F																											
B35	27	M									1.2	1.0	93	3															
B36	17	F											36	3															
B37	22	F											29	0.5															
B38	16	F											96	1															
B39	49	M	6.2	0.9	1						402	2																	
B40			4.8	1.2																									
													33	0.5															
B42	56	F											30	0.5															
B43	16	F											90	1															
B44	30	F	11.6	2.1	1		3.8	3.2			92	2																	
B45	44	M	5.4	0.5	1		4.6	1.6			168	2																	
B46			1.9	0.4																									
B47	29	F									107	2																	
B48	37	M																											
B49	45	M	8.9	2.1	1						56	2																	
B50	47	F									18	0.5																	

500074

INDIVIDUAL RADIOLOGICAL MONITORING DATA - BIKINI

(Urine Levels - amt/l; ^{137}Cs Body Burdens - 1974, nCi)

No.	Age	Sex	1973			1974			1975 (Mar.)			1975 (Oct.)			1976 (Mar.)			1976 (Oct.)			
			^{90}Sr pCi	^{137}Cs nCi	Yrs. Bik.	^{90}Sr pCi	^{137}Cs nCi	$^{137}\text{Cs}^{**}$ nCi	Yrs. Bik.	^{90}Sr pCi	^{137}Cs nCi	Yrs. Bik.									
B51	33	M				3.0	0.6	222	2							6.8	2.8	4			
B52	19	M						77	1												
B53	46	M						156	1	0.5	0.5	2				2.3	1.4	3			
B54	51	M						72	5												
B55	56	M														2.8	1.1	3			
B56	54	F														5.1	3.2	6			
B57	24	M														3.0	1.0	3			
B58	22	F														7.6	0.9	4			
B59	20	M														4.3	1.9	3			
B60	24	M														1.5	1.6	2			
B61	30	M														5.5	1.1	3			

** Gamma Spectrographic analyses - body burdens.

5009765

Mean Cesium-137 Levels Obtained by Whole Body Counting - 1974

	Male			Female		
	No.	nCi	nCi/kg body wt.*	No.	nCi	nCi/kg body wt.*
Bikini	8	128	1.84 (0.43-5.11)	13	73	1.15 (0.22-3.26)
Utirik	9	262	4.05 (2.64-6.84)	13	133	2.13 (0.96-3.85)
Rongelap	22	475	7.76 (4.37-16.3)	24	304	5.13 (2.71-13.46)
BNL med. team	4		2.93 0.0352(0.0134-.0791)			

* MPC 43 nCi/kg

Estimated Dose to Bone Marrow (mrem/yr)*

SOURCE	BIKINI	ENUE	RONGELAP	UTIRIK	DENVER	USA LONG ISLAND
Natural	80	80	80	80	325	190
Medical						
Dental	0	0	10	10	70	70
Contamination						
Gamma	165	7	20	7		
Internal	21	21	68	31		
TOTAL	266	108	178	128	395	260

* Dose on Marshall Islands based on personnel and environmental data.

009766
5009744

ANIMAL DATA
COCONUT CRABS

Island	Year	Part	nCi/kg	⁹⁰ Sr	¹³⁷ Cs	²³⁹⁻²⁴⁰ Pu
					nCi/kg	pCi/kg
Rongelap	1964	edible	4.0 - 9.0			
"	1969	whole	53.0		8.5	
"		"	34.0		7.0	
"	1972	"	56.0		13.0	
		"	28.6		5.0	
		"	39.0		6.1	
		"	31.0		6.8	
		"	23.0		5.1	
	1974	"	18.0		5.0	
		"	40.0		8.2	
		"	18.0		4.8	
	1975	{ edible	0.4		2.7	1.9
" Arbur Is.		skeleton	26.4		1.4	8.3
" Kabelle		{ edible	2.4		21.3	8.2
Island		skeleton	0.1		8.8	14.8
"		{ edible	0.8		14.6	5.1
		skeleton	24.1		5.0	2.2
Bikini		whole	23.3		11.8	1.5
	1970	"	24.8		14.8	0.07
	1971	"	132.0		11.4	
		"	412.0		8.6	
	1973	"	45.7		9.3	
	1974	muscle	16.0		380.0	18.0
		hepatopancreas	36.0		93.0	25.0
		skeleton	160.0		70.0	21.0
Enue	"				0.3	

5009767

ANIMAL DATA (1974)

OTHER THAN CRABS

Atoll	Sample	^{90}Sr pCi/kg*	^{137}Cs nCi/kg*	$^{239-240}\text{Pu}$ pCi/kg*
Pig				
Rongelap (Kabelle)	meat	10	11.0	
	organs	5	3.4	2.2
	bones	11,000	6.0	7.0
" Rongelap Island	Pig			
	meat	1.5	2.0	
	bones	1,600	0.9	0.4
"	chicken			
	meat	4	0.7	
	bones	480	0.6	0.8
"	meat	36	1.1	
	bones	3,500	0.6	0.6
Bikini Island	chicken			
	meat	8	0.7	
	bones	750	0.8	0.9
Rongelap Island	Rat 1	350	3.6	
	" 2	280	4.2	
	" 3	540	4.3	
Bikini Is.	Rat 1	6,000	8.1	6.7
	Fish	60		4.0
	(dry wt.)	240		20.0
		180		45.0
	" surgeon muscle			<.001 dpm/g

* Wet weight except dry wt. for fish.

5009768

TABLE 4

 RADIOCHEMICAL ANALYSES OF WELL WATER (pCi/liter) AND SLUDGE FROM WATER
 CISTERNS (pCi/g. dried sludge) FROM BIKINI

Year	Sample	Vol., ml	^{90}Sr *	^{137}Cs **	^3H	$^{239-240}\text{Pu}$ ***
1971	"good well"	1830	6.0 \pm 17%	600	\pm 1% 770 \pm 40%	0.04 \pm 25%
	"bad well"	1830	25 \pm 3%	850	\pm 1% 1040 \pm 30%	0.05 \pm 20%
	"good well" (closed)	1810	103 \pm 2%	1044	\pm 1%	0.058 \pm 15%
	"good well" (opened)	1980	125 \pm 3%	818	\pm 1%	5.76 \pm 6%
	drinking water (camp area)	3580	0.46 \pm 4%	1.53 \pm 8%		0.004 \pm 100%
1972	well water	1000	15.4 \pm 9%	800	\pm 1%	
	drinking water	1960	0.61 \pm 6%	1.8	\pm 8%	
1973	new well	60	52	600		0.38 \pm 40%
	B-1 well	225	11	724		0.08 \pm 50%
1975	Sludge from water cistern (6 samples)					2.91-4.02 \bar{x} 3.06

* MPC 4×10^3 pCi/l** MPC 2×10^5 pCi/l*** MPC 3×10^5 pCi/l

5009769

		<u>PLANTS</u>							
Year	Atoll/Island	Tissue	⁹⁰ Sr Range	137 Cs Avg.	Range	Avg.	239 Pu Range	241 Am Range	Avg.
<u>1. Bread Fruit (pCi/g dry)</u>									
74									
75	Bikini	leaves	251 - 446	332	29 - 95	53	.043-.148	.10	
74	Rongelap	leaves	23		27				
75	Wotho	leaves	.21		.35				
		fruit	<.08		1.7				
75	Bigej			- no data -					
<u>2. Pandanus (pCi/g dry)</u>									
74									
75	Bikini	leaves	41 - 402	170	37 - 1035	508	.031-.20	.087	
		fruit	34 - 255	138	3524 - 3665	3591	.001-.01	.005	
74	Rongelap	leaves	11		13				
		fruit		- no data -					
75	Wotho	leaves		- no data -					
75	Bigej	leaves	<.10		.92				
		fruit	.15		.55 - 1		.78		
<u>3. Coconut (pCi/g dry)</u>									
74									
75	Bikini	leaves	5.2 - 35	15.7	58 - 649	154	.018-.96	.205	
74	Rongelap	leaves	2.1		4.2				
75	Bigej	meat	<.15		1.2-2.3		1.75		
75	Wotho	leaves	.10		.70				
<u>4. Messerschmidia (pCi/g dry)</u>									
74	Bikini	leaves	235		305				
75		leaves	15 - 384	102			.07 -.985	.478	
74	Rongelap	leaves		- no data -					
75	Bigej	leaves	.16		2.3				
75	Wotho	leaves		- no data -					

5009110

Year	Atoll/Island	Tissue	<u>PLANTS</u>						
			⁹⁰ Sr Range	Avg.	¹³⁷ Cs Range	Avg.	²³⁹ Pu Range	Avg.	²⁴¹ Am Range
<u>5. Scaevola (pCi/g dry)</u>									
74	Bikini	leaves	33		110				
75		leaves	31 - 169	92			.07-.931	.252	
74	Rongelap	leaves	3.0		27				
75	Bigej	leaves	<.17		.95				
75	Wotho	leaves		- no data -					
<u>6. Papaya (pCi/g dry)</u>									
74	Bikini	fruit	81		1050		<.002		
75		fruit	74 - 79	76			.001-.009	.005	
75	Wotho	fruit	<.15		0.32				
		rind	<.14		15				
		seeds	<.14		12				
<u>7. Squash (pCi/g dry)</u>									
74	Bikini/	fruit	10		794				
	Bikini	fruit	5				.003		
<u>8. Banana (pCi/g dry)</u>									
74	Bikini	fruit	7.9		16				
75		fruit	9.33				.002		
		skin	90				.18		
<u>9. Arrowroot (pCi/g dry)</u>									
75	Bikini	tubers					.239		

500977

SOIL, AIR AND HOUSE DUST

Atoll/Island	Year	<u>SOIL</u>									
		⁹⁰ Sr Range	pCi/g. dry Average	¹³⁷ Cs Range	pCi/g. dry Average	²³⁹ Pu Range	pCi/g. dry Average	*	²⁴¹ Am Range	pCi/g. dry Average	
Bikini/Bikini	1970	-	-	27 - 955	199	7 - 140	43	5 - 45	17.5		
	1972	-	-	-	-	2.5 - 132	21	.85 - 8.1	3.8		
	1974	5.8 - 370	130.6	-	-	.44 - 9.3	3.				
	1975	18 - 432	127	14.9 - 250	24.5	.55 - 73	23	.24 - 17	6.5		
		14 - 578	202	-	-	-	-	-	-		
Kwajalein/Bigej	1975	.4	-	-	-	-	-	-	-		
Wotho/Wotho	1974	.3	-	-	-	-	-	-	-		
	1975	-	-	1.1	-	-	-	.08	-		

* MPC for Pu established by Eniwetok Task Group is 40 pCi/g.

AIR SAMPLES

(pCi/M³)

Bikini/Bikini	1972	0.06×10^{-4}	$0.12 \times 10^{-4}^*$
		to 0.2×10^{-4}	

* MPC is 6×10^{-3} pCi/M³ (ICRP, Pub. 2, 1959).

HOUSE DUST

(pCi/g)

Bikini/Bikini	1975	1.97 - 3.3	2.2
---------------	------	------------	-----

BODY BURDENS - STRONTIUM - 90
 (BASED ON RADIOCHEMICAL URINE ANALYSES)




Fig. 1

BODY BURDEN GAMMA EMMITTERS -
 WHOLE BODY GAMMA SPECTROSCOPY




Fig. 2

5004113