

RICHARD COLE

403060

Revised 3-67

FALLOUT CONTAMINATION OF FOOD AND WATER

by

Jack C. Greene
Director Postattack Research Division
of the
OFFICE OF CIVIL DEFENSE
Washington, D. C.

Based on a Paper Presented At the
Scientific Working Party Meeting
The Civil Defense Committee, North Atlantic Council
of the
North Atlantic Treaty Organization, Permanent Headquarters
Paris, France

May 1966



It is essential that protection against the external gamma radiation be prepared preattack. A policy depending on improvization in the post-attack period may be very dangerous. There probably would be insufficient time. On the contrary, if indeed the internal emitter problem should turn out to be very serious, there would be time available to take protective action. Obviously, the more we know about ways of reducing this hazard, and the better our plans are for coping with it, the better off we would be if we should have to use them.

It is the objective of this paper to discuss problems of fallout contamination in food and water as we view them today in the civil defense context. After outlining certain pertinent background experience that has led to our current understanding of the biological and physiological effects of the more dangerous radionuclides, I will mention our experience in the development of explicit standards and measurement techniques and equipment, and why we now consider such an approach to be unworkable and unnecessary. Then I will touch on the Office of Civil Defense research program, emphasizing our procedures for predicting the contamination levels likely to be associated with various types of nuclear wars, including quantitative estimates of maximum internal doses associated with certain hypothetical wars. Finally, I will mention the countermeasures that seem to merit consideration.

Probably the first human death from acute radiation poisoning was reported at a Berlin Medical Society Meeting in 1912.¹ A woman who had been treated for arthritis over a period of 16 days with injections of thorium-X, a short-lived isotope of radium, died within a month. The attending pathologist clearly recognized the cause of death. He reported, ". . . one cannot doubt for a minute that we have here a case of death caused by mesothorium." But the danger signals went largely unheeded.

The story of the painters of luminous watch dials has become a classic in the annals of industrial hygiene.^{1,2,3} Girls, employed as dial painters during the period 1914 to 1925, achieved a fine point on paint-laden brushes by shaping them with their lips. Minute amounts of radioactive material thus were ingested. In 1922, -23, -24, nine of these girls died with severe and unexplained anemia, and destructive lesions of jawbone and mouth. Many others have died of radium poisoning over the years, although some are still alive.

Also during the early part of the century, there was a fad of administering radium by injection and by mouth for ailments ranging from bad colds and broken bones, to insanity and old age. One Chicago physician gave radium to more than a thousand patients -- probably several thousand -- his own children included. Ironically, these

victims of radium poisoning tended to come from the higher-income class since the radium treatments were not inexpensive.

During this period, one group of mental patients received known amounts of radium chloride intravenously for therapeutic purposes. Follow-up studies of these patients, involving repeated and comprehensive observations, have yielded extensive data on the detection, distribution, retention, and injury sequelae of the medication.

Continuing studies of the radium poisoning cases are being carried on by Dr. Robley Evans and his co-workers at the Massachusetts Institute of Technology, Cambridge, Massachusetts; and at the AEC's Argonne National Laboratory in Chicago. Some additional, but more limited studies are being carried on elsewhere in the United States.

In 1941, standards for maximum permissible body burdens for radium for man were established by an Advisory Committee* to the U. S. National Bureau of Standards.⁴ The maximum body burden was set at 0.1 microcurie -- the value that is still in use today and the value adopted by the International Commission on Radiological Protection.⁵

Additional information about the effects in man of internally-deposited radionuclides comes from experience with the fission-product radioiodine produced during nuclear weapons tests. The case of the Rongelap Atoll natives is well known. Accidentally they were exposed to the

* This Committee is now called the National Council on Radiation Protection and Measurements and frequently is referred to as NCRP.



"exposed at less than 10 years of age, had nodules (68%). In addition, there were two boys with hypothyroidism in that group who had previously shown considerable growth retardation. These boys have improved on thyroid hormone therapy instituted six months ago. Six Rongelap people with nodules have been operated upon, 5 children in whom the nodules proved to be benign and one woman who had cancer of the thyroid. The latter case is now doing well.

"Of interest was the absence of thyroid abnormalities in the 60 Utirik children examined who were in the same age range of the high-incidence group of Rongelap-exposed children, but who had received considerably less exposure.

"Five of the Rongelap thyroid cases will be brought to the Medical Research Center at Brookhaven National Laboratory in May 1966, for further evaluation and possible surgery."⁸

There seems little question that the radioiodine entered the bodies of the Rongelap natives as a result of their drinking water from open cisterns.

Another possible source of information on effects of radioiodine in man is experience within the continental United States resulting from fallout from atmospheric weapons tests in Nevada during the period 1951 to 1962.



Although most of the local fallout was confined to the tests site, occasionally some extended into inhabited areas. The estimates are that the total doses to people within these areas were low. Dr. Gordon Dunning of the U. S. Atomic Energy Commission has estimated that the highest external exposure for any one person was 13.5 roentgens and to any community was 6 roentgens.⁹ Most of the doses were far lower than these values. Some fallout radionuclides, however, were deposited on pasture land, and they could have been ingested as a result of drinking fresh milk from the grazing cattle. Children were at greatest risk because of the smaller size of their thyroids, and because children usually drank more milk than adults. During the Fall of 1965, physicians examined some 2,000 of the Utah children from areas where the fallout occurred and, as a control, some 1400 children from Arizona where there was little or no fallout. Preliminary findings indicated thyroid abnormalities, as evidenced by the presence of nodules, in appreciably more of the Utah children than in those of Arizona.^{10,11}

In March of this year, a further report on the results of the Utah and Arizona study was released by the U. S. Public Health Service. This report stated that no malignant growths of the thyroid gland were found in any of the children. The study revealed, however, a number of cases of thyroiditis, an inflammation of the thyroid that may produce



The next phase of this study is now underway -- May 1966. Thyroid specialists are again examining those children originally thought to have thyroid abnormalities. I would assume the results will be available in the near future.

There seems to be firm evidence that weapon-produced radioiodine can produce physiological effects. In the case of the Rongelap incident, the thyroid damage was produced by the fallout from a single nuclear explosion. In Nevada, if indeed the thyroid abnormalities are found to be attributable to radioiodine (and as I just mentioned, no such causal link has been established), the fallout from a number of test detonations may have contributed, and if the radioiodine should be implicated, there seems little question that its route of entry was through fresh milk. Drinking water supplies predominantly are from wells.

During the Atomic Energy Project of World War II, large research programs in radiobiology were undertaken; and through the use of animal experiments, a great deal of knowledge has been generated on various internally-deposited radionuclides. But the basic source of information on the bone seekers (and this includes strontium-90) stems from the early radium experience.

The animal experiments have shown that certain of the bone-seeking nuclides produce greater damage to the bone than radium for the same



dosage.⁵ This can be attributed to several factors, including greater sensitivity for portions of the bone where the nuclide is deposited, and greater importance of damaged tissue. These factors are taken into account in establishing maximum permissible values.

Determinations of maximum permissible values for body burdens of the various radionuclides are based on two somewhat differing criteria. One criterion, as I have just discussed, is for the bone-seekers; the second, for the other radionuclides, is based on limiting the weekly dose to the organ where the nuclide concentrates to a value commensurate with the limitation for whole-body external exposure. Calculations have been made for the gamut of nuclear-weapon-produced radionuclides and the results may be found in the NRRP report "Maximum Permissible Amounts of Radioisotopes in the Human Body and Maximum Permissible Concentrations in Air and Water," U.S. Department of Commerce, National Bureau of Standards Handbook 53, March 20, 1953. This report was superseded by the NCRP Handbook No. 69 published in June 1959 and entitled "Maximum Permissible Body Burdens and Maximum Permissible Concentrations of Radionuclides in Air and Water for Occupational Exposure." These calculations also are reflected in the "Report of ICRP Committee on Permissible Dose of Internal Radiation (1959)" published in the 1960 issue of Health Physics.

The first official Civil Defense guidance on levels of radioactivity in food and water to be permitted under emergency conditions in war-time was developed in 1950 by Dr. William F. Bale of the University of Rochester while on temporary assignment to the Division of Biology and Medicine of the U.S. Atomic Energy Commission. I should like to quote from Dr. Bale's report.¹⁴

"It is probably not generally realized that water, and also food, can be very appreciably radioactive as measured by many portable radiation monitoring instruments now available and still be perfectly acceptable for human use under emergency conditions. The danger at the present time is probably greater that the presence of small amounts of radioactivity will lead to unwarranted shutting off of a municipal water supply or to a proclamation forbidding its use for drinking purposes thus causing an acute exacerbation of emergency conditions that may exist, rather than that consumption of contaminated water will cause significant damage to a military organization or a civilian municipality."

At about the same time, Mr. Adrian Dahl of the University of Rochester, and later Dr. Edwin P. Laug of the U.S. Food and Drug Administration prepared comparison standards to be used with a conventional-type Geiger counter survey meter for evaluating the degree of contamination of a sample of food and water. In 1952, the FCDA published two bulletins; one provided guidance on the levels of contamination acceptable under emergency conditions¹⁵ (based on Dr. Bale's work); and the other described methods for making measurements, and

The level was 1/3 this amount if the consumption period was assumed to last for 30 days.

This guidance was accepted as doctrine by U.S. Civil Defense organizations over the next several years. The count-rate produced by a few cc's of sample containing 9×10^{-2} microcuries per cc is easily measured by an ordinary Geiger counter instrument. The Geiger tube probe with the beta shield open was placed over a compound in the lid of a standard 4-ounce ointment tin containing an amount of U-238 or radium that empirically had been adjusted to give the count rate desired. The unknown water or food sample was placed in the bottom section of the tin, and a comparison of the two readings indicated whether the radioactivity of the sample was above or below the acceptable level. These comparison standards were purchased by OCD and distributed so as to be available for emergency use.

Today both the standards and the measurement techniques are no longer used by U.S. Civil Defense for the following reasons. First, the fallout is now known to be a principal hazard of a nuclear war. Studies of hypothetical nuclear attacks show that fallout could seriously affect very large areas of the country. Thus, adherence to the conservatism reflected in the earlier calculations could unnecessarily deny the use of badly-needed food and water. Second, because of high backgrounds expected from gamma radiation sources outside a protected

location, and from the almost inevitable beta contamination even within a shielded area, the measurement equipment probably would not operate because the background levels would drive it off scale.


Without definitive emergency standards, and without food- and water-monitoring equipment, what is to be done?

We believe that the internal-emitter hazard is so small compared to the external gamma radiation hazard that essentially it can be ignored, at least during the shelter stay-period. Later, as health standards return to more-nearly-normal peacetime criteria, evaluations of food and water based on radiochemical analyses will be necessary so that the specific nuclides can be identified, and precautions or countermeasures can be taken accordingly.

There are two common-sense procedures that should be followed in any fallout area: first, primarily because of the radioiodine problem, water from open reservoirs should be avoided if other sources are available; and second, during the first few weeks following the attack, children (especially babies), also because of the radioiodine hazard, should not be given milk from cattle that have grazed on contaminated pasture. However, in any situation where the requirement for food or water became stark, always it would be better to use any available source of supply than to deny it.

Figure 2

SCHEMATIC OUTLINE OF MODEL SYSTEMS FOR ESTIMATING RADIOLOGICAL EFFECTS



"soluble nuclide deposit." On the right, the WORLD-WIDE FALLOUT MODEL, taking into account both offensive strikes and counterattacks, produces data on "soluble nuclide deposit up to planting time." This data and information on "soil parameters" feeds into the ROOT UPTAKE MODEL, as does the data on "soluble nuclide deposit" from the LOCAL FALLOUT MODEL.

Similar considerations and appropriate mixing of outputs from the local- and world-wide fallout models apply to the FOLIAR CONTAMINATION MODEL and the WATER CONTAMINATION MODEL. Account is taken of those plants which would be killed by external gamma radiation exposure. Eventually data are produced on "nuclide concentrations in water, pastures, and food crops at harvest." The outputs split into two parts: the first goes through the ANIMAL DIET MODEL and ASSIMILATION MODEL, eventuating in "nuclide concentration in meat, milk, and eggs, " the other path takes into account the PROCESSING MODELS, TRANSPORTATION MODELS, and LOCAL DISTRIBUTION MODELS and, after considering availability of stockpiled supplies, the output is fed through the DIET MODEL and HUMAN ASSIMILATION MODEL, arriving at "absorbed dose" data which then may be translated into the "occurrence probabilities of late effects."

The analysis by Mr. Lee dealt with the effects of drinking filtered but otherwise unprocessed surface-source waters that were contaminated by fallout deposited directly into the surface sources. Two hypothetical attacks were studied: first, a heavy, mixed attack against city and counterforce targets with a total yield of 12,000 MT; and second, a city-avoidance, medium counterforce attack with a total yield of 7,000 MT. In both attacks the weapons were assumed to be surface burst, and the yields were assumed to be 50% from fission and 50% from fusion. Because only the soluble radionuclide concentrations were of interest, computer print-outs listed the maximum soluble concentrations of Sr^{89} and Sr^{90} , Ru^{106} , I^{131} , Ce^{137} , and Ba^{140} . As a measure of the radiological situation, the computer print-out also listed the external gamma radiation dose rates. It was assumed that everyone drank 1 liter of water per day. The absorbed doses for various organs derived from ingestion of the most contaminated water (this happened to be St. Louis, Missouri, which was assumed to be a target city) were calculated. It was estimated that no serious biological effect in adult humans would be expected from the consumption of the most contaminated water (even without the benefit of decontamination by normal water-treatment methods). The probable exception of this conclusion for the entire population, is for the thyroid doses to young children, where continued consumption

4. The contamination of surface waters by run-off from contaminated land areas probably would not exceed the initial contamination from direct deposit of fallout in exposed water sources.

5. Water-softening and water-purification plants would provide sufficient decontamination of the water to reduce significantly the absorbed doses to body organs.

6. In the postattack environment, water-source contamination would not be as critical a problem as the distribution of water for the survivors in damaged cities.

The other study that I want to mention relates to the contamination of human food following nuclear attack. It is the one that was performed for OCD by Isotopes, Inc.

The following diagram (next page) illustrates the approach. Radioactivity produced in the attack is separated into three categories -- that injected into the stratosphere, that into the high troposphere, and that into the local atmosphere. Estimates of the amount and nature of local fallout were based on Dr. Miller's models. Fallout of stratospheric debris was obtained from studies performed for the Defense Atomic Support Agency under Project Stardust. Consideration then was given, using the local fallout data, to the determination of contamination of crops that would be standing at the time of attack. Future crop



COMPUTATION OF THE RADIATION DOSE TO MAN FROM THE INGESTION OF CONTAMINATED FOOD

PROJECT DIAGRAM


I. Attack Model
 Quantity
 Composition
 Geographical Distribution

II. Contamination of Environment
 Deposition of Fallout
 Ground Air Concentration C^{14}

III. Contamination of Foodstuffs
 Foliar Absorption
 Root Absorption
 Grazing

IV. Ingestion
 Diet Concentration
 Biological Half-Life
 Critical Organ Concentration

V. Radiation Dose



rationing in order to provide the least-contaminated food to the young. If only a highly-contaminated supply were available, the uptake by the thyroid of I^{131} could be blocked through administration of stable iodine.

In a study sponsored by the U.S. Public Health Service, 21 sodium iodide was given in increasing doses to groups of children having normal thyroid function. The size of the sodium-iodide dose was varied in proportion to the skin area of the children. Maximum suppression of iodine-131 uptake was achieved with 1.5 to 2.0 milligrams of iodide per square meter of body surface per day. It quickly rebounded when the iodide was discontinued.

Based upon this experience, the research team calculated that the minimum dose of iodide required for almost complete suppression of the uptake of radioactive iodine by the normal human thyroid was 1.5 to 2.0 milligrams per day per square meter of body surface. For an adult, this dose of iodide would be 3 - 4 mg per day; and for children, about 1 mg per day.

At these iodide doses, suppression of uptake of radioiodine would begin almost immediately, and by 24 hours a 50% reduction would be expected. Subsequently, a gradual decrease in uptake to a minimum of about 5% would occur in 4 - 6 weeks. Toxic effects of iodide from daily doses of this order of magnitude given over relatively short periods of time would not be expected.

Administration to provide the rationale for allocating such foods. Since the physiological effects of high body burdens of strontium, like radium, are not expected until years after ingestion, older people obviously could tolerate the most contaminated of the supplies. Therefore, plans for allocation on the basis of age would seem to be desirable.

In a postattack world, the least contaminated food could become relatively expensive, a result to be expected if only the normal supply-and-demand factors were operative and no governmental control exercised. Thus, the more affluent of the survivors would be the least affected, and the poorer, the most affected. The situation easily could be exacerbated because of an exaggerated fear of contamination, the situation that probably exists among much of the population now. Thus, because the utilization of strontium-bearing food in a postattack world has not only physical and biological elements, but sociological, psychological, and economic implications as well, it needs careful study and planning now.

If, in a postattack situation, strontium-90 turned out to be a much greater hazard than the calculations indicate, and the countermeasures proved far less effective than expected, the consequences, although no doubt catastrophic in the eyes of those directly affected, would not be catastrophic in the sense of jeopardizing survival of the society.

LIST OF REFERENCES CITED

1. Schubert, Jack and Ralph E. Lapp, Radiation: What It Is and How It Affects You, The Viking Press, New York, 1957.
2. Claus, Walter D. (Ed.), Radiation Biology and Medicine, prepared for the United States Atomic Energy Commission, Addison-Wesley Publishing Co., Inc., Reading, Massachusetts, 1958.
3. Evans, Robley D., "Radium and Mesothorium Poisoning and Dosimetry and Instrumentation Techniques in Applied Radioactivity," MIT-952-2, Massachusetts Institute of Technology, May 1965.
4. "Safe Handling of Radioactive Luminous Compounds," Handbook H 27, National Bureau of Standards, 1941.
5. "Report of ICRP Committee II on Permissible Dose for Internal Radiation (1959), with Bibliography for Biological, Mathematical, and Physical Data," Health Physics Vol. 3, June 1960, Pergamon Press, New York.
6. Cronkite, E.P., V.P. Bond, and C.L. Dunham (eds), "Some Effects of Ionizing Radiation on Human Beings," U.S. Atomic Energy Commission, TID-5358, July 1956.
7. Conard, Robert A., and Aroboti Hicking, "Medical Findings in Marshallese People Exposed to Fallout Radiation," Journal of American Medical Association, Vol. 192, No. 6, pp 457-459, May 10, 1965.
8. Private communication dated April 14, 1965 from Dr. V.P. Bond, Chairman, Medical Department, Brookhaven National Laboratory, Upton, Long Island, New York, based on material prepared by Dr. Robert A. Conard of the Brookhaven Medical Department.
9. "Fallout from Nuclear Weapons Tests," Hearings before the Special Subcommittee on Radiation of the Joint Committee on Atomic Energy, Congress of the United States, 86th Congress, May 5, 6, 7, and 8, 1959, Volume 3, U.S. Government Printing Office, Washington, D.C. 1959 (starting on page 2021).

10. Press Release HEW-H25, U. S. Department of Health, Education, and Welfare, Public Health Service, Washington, D. C., October 28, 1965.
11. "Health Officials Study Thyroid Symptoms of Children Near Nevada Atomic Test Site," Wall Street Journal, October 28, 1965.
12. Press Release HEW-J74, U. S. Department of Health, Education, and Welfare, Public Health Service, Washington, D. C., March 16, 1966.
13. "Radiation, Fallout in Utah," Time, Vol. 87, No. 12, pp 60, March 25, 1966.
14. Memorandum for Files, William F. Bale, Subject: "Levels of Radioactivity in Water and Food that Can be Permitted Under Emergency Conditions in Wartime," dated November 10, 1950.
15. "Permissible Emergency Levels of Radioactivity in Water and Food," Civil Defense Technical Bulletin, TB-11-8, Federal Civil Defense Administration, December 1952.
16. "Emergency Measurements of Radioactivity in Food and Water," Civil Defense Technical Bulletin, TB-11-9, Federal Civil Defense Administration, December 1952.
17. Miller, Carl F., "Fallout and Radiological Countermeasures, Volumes I and II," SRI Project IM 4021, Stanford Research Institute, January 1963.
18. Miller, Carl F., "Fallout Models and Radiological Countermeasure Evaluations," SRI Project MU-5116, Stanford Research Institute, May 1965.
19. Lee, Hong, "Vulnerability of Municipal Water Facilities to Radioactive Contamination from Nuclear Attacks," SRI Project IM-4536, Stanford Research Institute, March 1964.
20. Bensen, David W., et al., "Summary Report on Contamination of Human Food Following Nuclear Attack. Vol. 1. Food Contamination Model, and Vol. 2. Derivation of Inputs for the Sr⁹⁰ Food Contamination Model," Isotopes, Inc., Westwood, New Jersey, October 1965.

21. Saxena, K.M., E.M. Chapman, and C.V. Pryles, "Minimal Dosage of Iodine Required to Suppress Uptake of Iodine¹³¹ by Normal Thyroid," Science, Vol. 138, pp 430-31, October 1962.
22. Parsi, Edgardo and William B. Iaconelli, "Removal of Radioactive Material from Milk by Electrodialysis," Ionics, Inc., Cambridge, Massachusetts, February 7, 1964.
23. Baldi, E.J., and Burdett Heinemann, "Strontium-90 Removal Project," Final Report to Department of Agriculture, Producers Creamery Co., Springfield, Missouri, September 25, 1965.
24. Ayres, Robert W., "Environmental Effects of Nuclear Weapons, Volumes I, II, and III (Summary)," HI-518-RR, Hudson Institute, 1 December 1965.

ADDITIONAL REFERENCES

- Grune, Werner N. and Henry S. Atlas, "Evaluation of Fallout Contamination of Water Supplies," Merrimack College, North Andover, Massachusetts, 15 May 1965.
- Lansing, N. F. (compiled by), "The Role of Engineering in Nuclear Energy Development," Proceedings of the Third Annual Oak Ridge Summer Symposium, Technical Information Service, Oak Ridge, Tennessee, TID-5031, December 1951.
- Miller, Carl F., "Biological Availability and Uptake of Fission Products in Fallout," SRI Project IMU-4021, Stanford Research Institute, April 1962.
- Lane, William B., James D. Sartor, and Carl F. Miller, "Plant Uptake of Radioelements from Soil," SRI Project IM-4536, Stanford Research Institute, Menlo Park, California, March 1964.
- "Maximum Permissible Amounts of Radioisotopes in the Human Body and Maximum Permissible Concentrations in Air and Water," Handbook 52, U.S. Department of Commerce, National Bureau of Standards, March 20, 1953.
- Conard, Robert A., et al., "Medical Survey of the People of Rongelap and Utirik Islands Nine and Ten Years after Exposure to Fallout Radiation (March 1963 and March 1964)," BNL 908 (T-371), Brookhaven National Laboratories, May 1965.
- Knapp, H. A., "Iodine-131 in Fresh Milk and Human Thyroids Following a Single Deposition of Nuclear Test Fallout," Nature Vol. 202, No. 4932, pp 534-537, May 9, 1964.
- Knapp, H. A., "Average and Above-Average Doses to the Thyroids of Children in the United States from Radioiodine from Nuclear Weapons Tests," prepared as part of a report on Radiation Exposure in the United States from Nuclear Test Fallout, U.S. Atomic Energy Commission, Washington, D. C., August 6, 1962.

Bustad, L.K., Biology of Radioiodine, Proceedings of the Hanford Symposium on the Biology of Radioiodine, Pergamon Press (1964).

Federal Radiation Council Reports Nos. 1-6, Superintendent of Documents, U. S. Government Printing Office, Washington, D. C.

The Biological Effects of Atomic Radiation; (a) Summary Reports 1960, (b) A Report to the Public 1960, National Academy of Sciences, National Research Council, Washington, D. C.

Report of the United Nations Scientific Committee on the Effects of Atomic Radiation, General Assembly Official Records: Thirteenth Session, Supplement No. 17 (a/3838), New York 1958.

Report of the United Nations Scientific Committee on the Effects of Atomic Radiation, General Assembly Official Records: Seventeenth Session, Supplement No. 16 (a/5216), New York, 1962.

