

CRITERIA FOR ESTABLISHING SHORT TERM
PERMISSIBLE INGESTION OF FALLOUT MATERIAL

Gordon M. Dunning
Division of Biology and Medicine
Atomic Energy Commission
Washington, D. C.

The criteria for establishing permissible ingestion of radioactive fallout material under emergency conditions for several weeks following a nuclear detonation are dependent primarily on exposures to the,

- a. gastrointestinal tract from the gross fission product activity,
- b. thyroid from the isotopes of iodine and,
- c. bone, principally from Sr^{90} - Y^{90} , Sr^{89} , Ba^{140} - La^{140} .

I. Doses to the Gastrointestinal Tract

The following principal assumptions are used in calculating the doses to the gastrointestinal tract of adults:

- a. The calculations are based on the methods contained in reference one.
- b. The fallout material is 90 percent insoluble. (See IV. Discussion below).
- c. The activity decays according to the principle of (time)^{-1.2}.

US DOE ARCHIVES 326 U.S. ATOMIC ENERGY COMMISSION
RG _____
Collection <u>DOS McCraw</u>
Box <u>4</u> Job <u>1320</u>
Folder <u>MHS 14 6.4 Fallout-MISC</u> <u>1957-1958</u>

CONFIRMED TO BE UNCLASSIFIED
BY AUTHORITY OF DOE/OC

REVIEWED BY

DATE

Carl W. ... 12/21/87

134
MJA



- d. The energy delivered is all derived from the beta emissions, having a mean energy of 0.4 Mev when in the lower large intestine. (See Graph One)²
- e. The total daily consumption of food and water is 2200 grams or milliliters.

The method of calculation is according to the following equation:

$$\frac{(\text{Total number of disintegrations occurring in organ})(\text{Energy of emissions})(8.0 \times 10^{-9})}{(\text{Mass of Organ})} =$$

Dose (rads) (1)*

The number of disintegrations taking place in the organ may be calculated according to equation two:

$$\text{Total number of disintegrations} = 5A_a t_a^{1.2} [t_a^{-0.2} - t_b^{-0.2}] \quad (2)$$

Where: A_a = number of disintegrations per unit time at time "a" after detonation.

t_a = time "a" after detonation.

t_b = time "b" later than "a".

One of the more useful forms for the criteria would be in units of permissible concentrations at time of intake. This will somewhat complicate the calculations since there will be a decrease in activity as the material passes along the gastrointestinal tract. When such calculations are made according to the above assumptions and equations, it may be seen that the critical organ is the lower large intestine except for the first hours immediately following the detonation. (Table One shows the relative doses

* The rad is the unit of absorbed dose equal to 100 ergs per gram.

$$\frac{1.6 \times 10^{-6} (\text{ergs/Mev}) 0.5 (\text{proportion of total energy to gastrointestinal tract})}{100 (\text{ergs/gm-rad})} = 8.0 \times 10^{-9}$$

DOE ARCHIVES

135
120

[REDACTED]

to parts of the gastrointestinal tract as a function of time.) Therefore, Graph Two is based on the activity at time of ingestion to produce one rad of dose to the lower intestine.

For example, Graph Two shows that if about 48 microcuries are ingested on the 24th hour after detonation, the lower large intestine may receive one rad of radiation dose. This was calculated in the following manner.

Step 1. Determine the total number of disintegrations in the lower large intestine necessary to produce 1.0 rad.

From equation (1)

$$\frac{(\text{Number of disintegrations}) (0.4) 8.0 \times 10^{-9}}{150} = 1$$

$$\text{Number of disintegrations} = 4.7 \times 10^{10}$$

Step 2. Determine the activity at time of intake to produce 74.7×10^{10} disintegrations within the large intestine.

$$\frac{4.7 \times 10^{10}}{0.9} = 5.2 \times 10^{10} \text{ disintegrations intake required (assuming 10\% solubility).}$$

$$5.2 \times 10^{10} = (5) (A_{37}) (37^{1.2}) [37^{-0.2} - 55^{-0.2}] *$$

$$A_{37} \approx 3.7 \times 10^9 \text{ d/hr.}$$

$$A_{24} \approx 6.2 \times 10^9 \text{ d/hr.}$$

$$A_{24} \approx 47 \mu\text{c}$$

* If the time of intake is the 24th hour, then the start of irradiation of the lower large intestine is $24 + 13 = 37^{\text{th}}$ hour.

DOE ARCHIVES

Graph Two has been used in estimating radiation doses to the lower large intestine for prolonged periods of ingestion (Table Two). The following calculations are illustrative for the period of 24th to the 120th

[REDACTED]

hour (start of intake at the beginning of the 2nd day after detonation for a duration of four days).

Step 1. Determine the number of microcuries at time of ingestion to produce 1.0 rad to the lower large intestine.

From Graph Two take the mid point of intake period (72nd hour) → 31 μc . (This is obviously an approximation since the exact times of intake during the four-day period will be unknown).

Step 2. Determine the activity at time of intake.

From equation (2)

$$31 = 5 A_{24} 24^{1.2} [24^{-0.2} - 120^{-0.2}]$$

$$A_{24} \cong 0.94 \mu\text{c/hr}$$

Since there is assumed a 220 ml/day intake

$$0.94 \times \frac{24}{2200} \cong 0.010 \mu\text{c/ml or gm}$$

II. Doses to the Thyroid

The following principal assumptions are used in calculating the doses to the adult thyroid from intake of activity from fallout material:

- a. The percentages of the isotopes of iodine in mixed fission products are according to Hunter and Ballau.³
- b. Twenty percent of the ingested I^{131} reaches the thyroid.
- c. The mean energy is 0.22 Mev.
- d. The thyroid weight is 20 grams. (See IV. Discussion below)
- e. The percentages of shorter-lived isotopes of iodine that reach the thyroid and their doses are according to reference four.

DOE ARCHIVES

The method of calculation of doses to the thyroid is illustrated by computing that amount of intake of fission products at the 48th hour

to produce 1.0 rad.

Step 1. Determine the dose rate on the day of intake of I^{131} to produce 1.0 rad to the thyroid.

$$D = \frac{R}{\lambda e} \text{ Where: } D = \text{dose (1.0 rad)}$$

$$R = \text{dose rate on initial day}$$

$$\lambda e = \text{effective decay constant (radiological and biological)}$$

$$1.0 = \frac{R}{0.09}$$

$$R = 0.09 \text{ rads/day}$$

Step 2. Determine the number of microcuries of I^{131} to produce 0.09 rad/day

$$\frac{X(\mu c)(2.2 \times 10^6)(60 \times 24)(1.6 \times 10^{-6})(0.22)}{(100)(20)} = 0.09$$

$$X = 0.16 \mu c \text{ to thyroid}$$

$$\text{or}$$

$$(0.16)(5) = 0.80 \mu c I^{131} \text{ ingested}$$

Step 3. Determine relative doses from I^{131} and I^{short} according to Graph Three.⁴

At 48th hour, the relative contribution to total dose from I^{131} and I^{short} is about $\frac{1}{1}$.

Therefore, ingestion of 0.4 $\mu c I^{131}$ (equivalent) at 48th hour will produce 1.0 rads to thyroid.

Step 4. Determine the number of microcuries of fission products required to yield the required I^{131} activity. At 48th hour, I^{131} constitutes about 2.35% of total activity.

Therefore,

DOE ARCHIVES

$$\frac{0.4}{0.023} \approx 17 \mu c \text{ of fission products.}$$

[REDACTED]

Graph Four shows the number of microcuries of fission products ingested at times after detonation to produce 1.0 rad to the thyroid.

III. Doses to the Bones

The three principal bone-seeking isotopes of concern are Sr^{90} - Y^{90} , Sr^{89} , and Ba^{140} - La^{140} . Evaluation of these may be made in terms of amount deposited in the bones versus maximum permissible body burdens, or in rads of dose that they deliver after deposition. Since values for maximum permissible body burdens are based on the concept that these will be maintained indefinitely in the body, they are not so valid for Sr^{89} and Ba^{140} - La^{140} when considering short periods of emergency intake.

The following principal assumptions are used in calculating the doses to the bones of adults:

a. The percentages of the isotopes of Sr^{90} - Y^{90} , Sr^{89} , and Ba^{140} - La^{140} in mixed fission products are according to Hunter and Ballau.³

b. The percentages of intake of these isotopes that are deposited in the bones, the energies of emissions, and their effective half lives are according to reference five - except for Sr^{90} where a 27.7 year radiological half life is used here.

c. The mass of the bones is 7,000 grams.

DOE ARCHIVES

The method of calculation of doses to the bones is illustrated by computing the dose from Sr^{89} from the intake of 27 microcuries (See IV Discussion below) of mixed fission products on the 120th hour. Similar calculations were made for Sr^{90} - Y^{90} and Ba^{140} - La^{140} and then the three doses

[REDACTED]

were added for each intake of fallout material.

Step 1. Determine the Sr⁸⁹ to reach the bone.

According to reference 4:

The Sr⁸⁹ content in mixed fission products on the 120th hour is 1.6%.

According to reference 5:

The intake of Sr⁸⁹ to reach to the bones is 25%.

Therefore:

$$(27) (0.016)(0.25) = 0.108, \text{ to the bone.}$$

Step 2. Determine the dose rate to the bones.

With an assumed effective energy of 0.55 Mev (reference 5).

$$\frac{(0.108)(2.2 \times 10^6)(60 \times 24)(1.6 \times 10^6)(0.55)}{(100)(7,000)} = 4.3 \times 10^{-4} \text{ rads/day}$$

or 0.43 millirads/day

Step 3. Determine total dose.

$$D \text{ total} = \frac{R}{\lambda e} \text{ where: } R = \text{initial dose rate} \\ \lambda e = \text{effective decay constant}$$

$$D \text{ total} = \frac{0.43}{0.0133} \approx 32 \text{ millirads}$$

IV. Discussion

A. Solubility

DOE ARCHIVES

The solubility of fallout material varies, depending among other factors upon the surface over which the detonation occurred. The fallout material collected in soil samples at the Nevada Test Site has been quite insoluble, i.e. only a few percent in distilled water and roughly 20-30 percent in 0.1 N HCl. However, it would be expected that the activity actually present in

140
~~140~~

[REDACTED]

drinking water supplies would be principally in soluble form. The water collected from a well and a cistern on the Island of Rongelap (Table Three) about 21 months after the March 1, 1954 fallout, was found to have about 80 percent of the activity in the filtrate, but there was an undetermined amount that settled to the bottom. Other data suggest the material to have been about 10-20 percent soluble in water.

In the event contaminated food is ingested it is possible that the total activity--soluble and insoluble--may find its way into the gastrointestinal tract since at times immediately following a fallout most of this activity probably would come from the surface contamination rather than the soil-plant-animal cycle. There may then follow some solubilizing in the acid stomach with subsequent removal from the tract before reaching the lower large intestine.

It is assumed for these calculations that (a) 90% of the fallout material is insoluble when computing doses to the gastrointestinal tract, and (b) that the isotopes of iodine, strontium, and barium are all soluble when computing doses to the thyroid and to the bones. These assumptions are probably conservative, i.e. they may overestimate somewhat the radiation exposures.

B. Biological Significance.

DOE ARCHIVES

After the estimation of radiation doses by any procedure the final step is an evaluation in terms of biological effects both for short and long terms.

0 [REDACTED]

1. Gastrointestinal Tract

There have been few experiments where the gastrointestinal tract has been exposed in a manner similar to the one assumed here. In one experiment currently underway at the Medical Center of the Oak Ridge Institute of Nuclear Studies, the dosimetry is being studied in dogs using Yttrium-90. The preliminary data are in reasonable agreement with the model proposed in reference one.⁶

In another experiment,⁷ rats were fed 1.0 to 6.0 millicuries of Yttrium-90 in a single feeding. Four of the 33 animals died of adenocarcinoma of the colon and additional animals died with acute and chronic ulceration of the colon. A second group of rats was given 0.46, 0.20, or 0.06 mc of Y^{91} per feeding over a period of three months with total accumulated amounts of 31.2, 15.6 and 4.68 mc respectively. Six of the eight animals at the two higher levels died with carcinoma of the colon and no malignancies were observed at the lowest level. The authors made no estimate of radiation doses.

DOE ARCHIVES

In another experiment,⁸ rats were kept alive by the use of parabiosis or para-aminopropio phenone either pre or post whole-body irradiation of 700-1000 roentgens. Four of the 21 rats developed tumors along the gastrointestinal tract (one each jejunum, ileum, duodenum, and colon), with four additional animals showing tumors in other organs. However, in comparing gastrointestinal versus whole-body irradiation, the question has been raised as to a possible indirect carcinogenic action in the latter case.⁹ By using fast neutrons, lesser doses have been shown to produce an appreciable percentage of intestinal carcinomas in mice, but this is not so relevant

142
~~157~~

[REDACTED]

to the present discussion of beta exposure.¹⁰

One summarizing statement of the short-term effects stated, "--- though the gastrointestinal tract is one of the sensitive systems to ionizing radiation, it also has a most remarkable regenerative and reparative capacity. It takes doses of well over a thousand roentgens to damage the gut permanently in most mammals studied, and it is capable of rapid, dramatic recovery of anatomical and functional integrity with doses in the lethal range."¹¹ Evaluating the data from dogs exposed to whole-body X-radiation the authors said, "--- it is suggested that doses of approximately 1,100 to 1,500 r may represent the upper limit of the possible efficacy of supportive measures in the treatment of the syndrome of acute radiation injury. With greater doses the damage to the intestinal mucosa appears irreparable and of an extent incompatible with life."¹² At the same time, it has been repeatedly indicated that the irradiation of the gastrointestinal tract plays a major role in gross whole-body effects associated with radiation syndrome.^{11, 12, 13, 14, 15, 16, 17, 18, 19, 20} In fact one author¹³ summarizes several experimental findings, "In producing acute intestinal radiation death, irradiation of any major portion of the exteriorized small intestine alone is almost equivalent to whole-body irradiation---."

Graph Five suggests the relative doses to the parts of the gastrointestinal tract, from ingestion of fallout material. The available experimental data does not permit a conclusive statement as to whole-body effects to be expected from such ratios of exposures. Most of these experiments are related to the criterion of death, but they do suggest

[REDACTED]

that the major contributory factor to such effects such as nausea and vomiting associated with whole-body exposures of 100-200 roentgens, may be the result of the gastrointestinal reaction. Possibly a few hundred rads to the lower large intestine together with the concomitant lesser exposures to the upper large intestine, the small intestine and the stomach (according to Graph Five) may be in the range where radiation sickness might occur.

2. Thyroid

The study and treatment of disorders of the thyroid gland with radioiodine has led to considerable information on doses and their effects to this organ. (Only a partial list of references is noted.)^{21,22, 23, 24, 25} Whereas these treatments have been principally with abnormal thyroids, much of the information may be extrapolated to normal thyroids for the purposes of this discussion. In addition there are other data based on normal thyroids in patients suffering such ailments as congestive heart failure.²⁶

The picture that is clearly presented is that of the relative insensitiveness of the adult human thyroid to radiation. For example, Freedberg, Kurland, and Herman,²⁶ report, "---Seven days after administration of 17 and 20 millicuries of I¹³¹, which delivered 14,500 and 31,000 rep, respectively, to the thyroid gland, no histologic changes were noted which could be attributed to I¹³¹. --- Fourteen and twenty-four days, respectively, after administration of 59 and 26 millicuries of I¹³¹, marked central destruction of the thyroid gland was noted.---" Since the first two patients expired seven days after administration of the I¹³¹ from pulmonary edema, it does not eliminate the possibility that the destructive

DOE ARCHIVES

[REDACTED]

changes might have appeared in the thyroid if these patients had survived. However, the evidence from other studies strongly indicate that if any pathological effects were to be noted in the thyroid after an exposure of some 10,000 reps they would be minimal. Likewise, the possibility of serious damage to other organs of the body, such as parathyroids and trachea which are simultaneously exposed to the I^{131} radiations, would be exceedingly small.

On long term effects, two summarizing statements may be made. "No thyroid neoplasm was found which could be attributed to I^{131} ,"²⁶ after doses to normal thyroids running into many tens of thousands of reps and after periods of observation up to more than eight hundred days. "In a series of over 400 patients treated with radioactive iodine at the Massachusetts General Hospital during the past ten years no known carcinoma of the thyroid attributable to this agent has developed. Definite answers to the question of carcinoma formation must await prolonged observation of treated patients."²³ Here the average treatment dose of I^{131} was 10 millicuries and of I^{130} 25 millicuries.

However, significantly lesser doses may be carcinogenic in children.²⁷

"---It has been suggested that the human thyroid is less radiosensitive than other tissues, such as bone, since after many years of treatment of Graves' disease with radioactive iodine, no cases of resulting carcinoma have been reported. The customary dosages of I^{131} in such cases yield at least 4000 rep to the gland. On the other hand, carcinoma of the thyroid found in children and young adults has almost invariably been preceded by x-ray treatment to the upper part of the body, in amounts such as to yield as little as 200 r to the infant thyroid. It has been estimated that less than 3% of such

DOE ARCHIVES

treated cases yield carcinoma; nevertheless, the data suggest that 200 r is a potentially carcinogenic dose to the infant thyroid. While the possibility exists that the carcinogenic action may be an indirect, hormonal one, it must still be recognized that this, like leukemia, is an instance of significant carcinogenesis by less than 1000 rep. It seems likely that the infant thyroid is unduly susceptible, but that the adult thyroid is not.---"28

Table Two indicates the amount of ingested fission product activity to produce one rad dose to the lower large intestine and Graph Five shows the relative doses to the gastrointestinal tract and the thyroid. It may be seen that ingestion of a given activity on the fourth and fifth days may result in nearly two and one-half times the dose to the thyroid as to the lower large intestine. For a continuous consumption of fallout material from the first hour to the 30th day the ratio of doses is about 1.7.

3. Bones

DOE ARCHIVES

It is recognized that the intake and deposition of strontium-89 and 90 are intimately associated with the calcium in the diet. Whereas it has been assumed here that a fixed percentage of the strontium intake is deposited in the bones (Reference Five). It is realized that this method involves uncertainties, as would the necessary assumptions to generalize for a wide variety of calcium--strontium ratios and intakes

to cover multiple categories. In situations where doses to the bones appear to be the critical criterion (such as later times after detonation than considered here), it would be necessary to make a more precise evaluation.

Unequal distribution of isotopes in the bones has been observed. Thus, the dotted line in Graph Five is included to suggest a possible larger dose to those regions.

Considerable data have been collected on radiation produced bone cancers. One summarizing statement that places this in proper perspective with the other factors discussed above is "--- Visible changes in the skeleton have been reported only after hundreds of rep were accumulated and tumors only after 1,500 or more."²⁹ When one examines Graph Five for relative doses, and reviews the data on doses versus effects to the gastrointestinal tract and possibly children's thyroids (Table Four), it would appear that exposure to the bones is not the critical factor for ingestion of fallout material under emergency conditions, for the first few weeks after a detonation.

4. Summary

DOE ARCHIVES

Table Four summarizes some possible biological effects from radiation exposures. Due to inherent uncertainties in such analyses together with expected wide biological variances among individuals, Table Four is intended only to suggest a generalized picture of doses versus effects.

The physical calculations of radiation doses made above were for

~~CONFIDENTIAL~~

adults. For equal intakes of radioactivity, children probably would receive higher exposures due to the smaller organ masses, and in the case of bones a greater deposition would be expected. Also, there is the possibility of tumor production in the thyroids of some children at relatively low radiation exposures. It would appear wise therefore to establish lower limits of intake of radioactivity for children.

C. Permissible Intake

The preceding discussion attempted to give estimates of radiation doses resulting from intake of fallout material, together with some possible biological effects. How much intake is actually permitted depends upon many factors including the essentialness of the food and water to sustaining life, and one's philosophy of acceptable biological risks and damage in the face of other possible hazards such as mass evacuation. Table Two and Graph Five give estimates of the amount of contamination in food and water to produce certain radiation doses to the critical organs. Table Four indicates possible biological effects from given doses. Using these references, command decisions may be made as to permitted intake of radioactivity. **DOE ARCHIVES**

Such evaluations as attempted here are necessary and valuable for planning purposes, but once the fallout occurs the emergency of the situation may preclude immediate analysis of the food and water supplies. Further, abstaining from ingestion of food and water because it might be contaminated could not be continued indefinitely. Therefore, the following three common sense rules are suggested:

████████████████████

1. Reduce the use of contaminated food and water to bare minimum until adequate monitoring can be done; use first any stored clear water and canned or covered foods; wash and scrub any contaminated foods and;

2. If the effects of lack of food and water become acute, then use whatever is available but in as limited quantities as possible. Whenever possible select what seems to be of the least likely contaminated water and/or foodstuffs; and

3. Since it is especially desirable to restrict the intake of radioactivity in children, give them first preference for food and water having the lowest degree of contamination.

In an area of heavy fallout one matter to consider is the relative hazards from the external gamma exposure versus internal doses from ingestion of the material. (Inhalation is thought to contribute only relative minor doses under the conditions discussed here). The best evidences on this point was the fallout that occurred on the Rongelapese in March 1954. Those in the highest exposure group received 175 r whole-body external gamma exposure yet their body burdens of internal emitters were relatively low (Table Five).³⁰ These and other data suggest that:

If the degree of contamination of an area is such that the external gamma exposure would permit normal and continuous occupancy after a fallout, the internal hazard would not deny it.

DOE ARCHIVES

This is based on such reasonable assumptions of (a) about 50% reduction of gamma exposure from out-of-doors doses afforded by living a part of each day in normal family dwellings, (b) washing

~~CONFIDENTIAL - EYES ONLY~~

and/or scrubbing contaminated foods, and (c) excluding areas where relatively little fallout occurred, but into which may be transported highly contaminated food and/or water. After longer periods of time during which the gamma dose rates in an originally highly contaminated area have decreased to acceptable levels, it probably would be necessary to evaluate the residual contamination for the bone seeking radioisotopes, especially strontium-90.

DOE ARCHIVES

215

R E F E R E N C E S

1. Maximum Permissible Concentration of Radioisotopes in Air and Water For Short Period Exposure. Morgan, K. Z., Snyder, W. S., and Ford, M. R. International Conference on the Peaceful Uses of Atomic Energy, July 8, 1955, Paper A/Cong. 8/P/79.
2. USNRDL-394. The Ratio of Lung Beta Dose to Whole Body Dose During Given Intervals After Atomic Detonation. (Confidential) Sondhaus, C. A., 1952.
3. Simultaneous Slow Neutron Fission of U²³⁵ Atoms I. Individual and Total Rates of Decay of Fission Products. Hunter, H. F. and Ballau, N. E., U. S. Naval Radiological Defense Laboratory, April 1949.
4. "Two Ways to Estimate Thyroid Dose From Radiiodine in Fallout," Dunning, Gordon M., Nucleonics Vol. 14, No. 2, February 1956.
5. Maximum Permissible Amounts of Radioisotopes in the Human Body and Maximum Permissible Concentration in Air and Water. Handbook 52, National Bureau of Standards, March 1953.
6. Private communication. Dr. Cyril Comar, Medical Division, Oak Ridge Institute of Nuclear Studies to Dr. Gordon Dunning. Work is sponsored by contract with Armed Forces Special Weapons Project, Department of Defense.
7. "Carcinoma of the Colon in Rats Following the Feeding of Radioactive Yttrium," Lisco, H., Brues, A. M., Finkel, M. P. and Grundhauser, W., Cancer Research 1947, Vol. 7 p. 721. DOE ARCHIVES
8. "Neoplasms in Rats Protected Against Lethal Doses of Irradiation by Parabiosis and Para-Aminopropio-phenone," Brecher, G., Cronkite, E. P., Peers, J. H., Journal of the National Cancer Institute, August 1953, Vol. 14, No. 1, p. 159-167.

~~CONFIDENTIAL~~

[REDACTED]

9. "Radiation as a Carcinogenic Agent," Brues, A. M. Radiation Research, November 1955, Vol. 3, No. 3, p. 272-280.
10. "Induction of Intestinal Carcinoma in the Mouse by Whole-Body Fast-Neutron Irradiation," Nowell, C., Cole, L. J., Ellis, M. E., Cancer Research, October 1956, Vol. 16 No. 9, p. 873-876.
11. Some Effects of Ionizing Radiation on the Physiology of the Gastrointestinal Tract. A Review. Lecture and Review Series No. 56-2. Conard, Robert, Naval Medical Research Institute, Bethesda, Maryland, March 1956.
12. "Lesions of the Alimentary Tract of Dogs Exposed to Whole Body X-Radiations of 300 to 3,000 R." Brecher, G., and Cronkite, E. P., American Journal of Pathology, 1951, Vol. 27, p. 676-677.
13. "The Nature of Intestinal Radiation Death," Quastler, H. Radiation Research, April 1956, Vol. 4, No. 4.
14. "Effects of Partial Shielding of Rat Intestine During X-Irradiation." Swift, M. N., Taketa, S. T., and Bond, V. P., Federation Procedures, 1954, Vol. 13, p. 523.
15. "Prevention of Intestinal Radiation Death by Removal of the Irradiated Intestine," Osborne, J. W., Radiation Research, June 1956, Vol. 4, No. 6.
16. "Sensitivity of Abdomen of Rat to X-irradiation", Bond, V. P., Swift, M. N., Allen, A. C. and Fishler, M. C., American Journal of Physiology, Vol. 161 1950, P. 323-330.
17. "Acute Intestinal Radiation Death. Studies on Roentgen Death in Mice, III." Quastler, H., Lanzl, E. F., Keller, M. E. and Osborne, J. W., American Journal of Physiology, February 1951, Vol. 164, No. 2, p. 546-556.

DOE ARCHIVES

[REDACTED]

- [REDACTED]
18. "Observations on Gastrointestinal Function After X-Ray and Thermal Column Exposures," Woodward, Kent T. and Rothermel, Samuel, Radiation Research October 1956, Vol. 5, P. 441-449.
 19. "Modification of Acute Intestinal Radiation Syndrome Through Shielding," Swift, M. N. and Taketa, S. T., American Journal of Physiology, April 1956 Vol. 185, No. 1, p. 85-91.
 20. "Lethal Effects of Intra-gastric Irradiation in the Dog." Littman, A., Fox, B. W., Schoolman, H. M., and Ivy, A. C., American Journal of Physiology, 1953, p. 347-351.
 21. "Radioiodine in the Study and Treatment of Thyroid Disease: A Review." Kelsey, M. P., Haines, S. F., Keating, F. R., The Journal of Clinical Endocrinology, Vol. 9, No. 2 February 1949, p. 171-210.
 22. Radioisotopes in Medicine, Andrews, G. A., Brucer, M. and Anderson, E. (Editors), September 1953. Superintendent of Documents, U. S. Government Printing Office, Washington 25, D. C. Chapter 18-27.
 23. "A Critical Analysis of the Quantitative I-131 Therapy of Thyrotoxicosis." Freeberg, A. S., Kurland, G. S., Chamovitz, D. L., Ureles, A. L. Journal of Clinical Endocrinology, Vol. 12 1952, p. 86-111.
 24. "Functional and Histologic Effects of Therapeutic Doses of Radioactive Iodine on Thyroid of Man." Dobyns, B. M., Vickery, A. L., Maloot, F., and Chapman, E. The Journal of Clinical Endocrinology and Metabolism. Vol. 13 No. 5, May 1953, p. 564.
 25. "Biological Hazards and Toxicity of Radioactive Isotopes", Brues, A. M., The Journal of Clinical Investigation. Nov. 1949, Vol. XXVIII, No. 6, p. 1286-1296.
- [REDACTED]

DOE ARCHIVES

~~CONFIDENTIAL~~

26. "The Pathologic Effects of I-131 on the Normal Thyroid of Man". Freeberg, A. S., Kurland, G. S. and Blumgart, H. L., The Journal of Clinical Endocrinology and Metabolism, Vol. 12, 1952, p. 1315-1348.
28. Pathologic Effects of Atomic Radiation. National Academy of Sciences - National Research Council, 1956, Washington, D. C.
27. "Association of Irradiation with Cancer of the Thyroid in Children and Adolescents," Clark, D. E., The Journal of the American Medical Association, Nov. 1955, Vol. 159, p. 1007-1009.
29. The Biological Effects of Atomic Radiation - Summary Reports. National Academy of Sciences -- National Research Council, 1956, Washington, D. C.
30. Some Effects of Ionizing Radiation on Human Beings. Cronkite, E. P., Bond, V. P., and Dunham, C. L. (Editors). Superintendent of Documents U. S. Government Printing Office, Washington 25, D. C.

DOE ARCHIVES

~~CONFIDENTIAL~~

A C K N O W L E D G M E N T

Acknowledgement is gratefully made for the kind assistance in reviewing the manuscript and offering many helpful suggestions: Dr. Charles L. Dunham, Director, Division of Biology and Medicine, AEC, Washington, D. C.; Dr. Forrest Western, Health Physicist, AEC, Washington, D. C.; Dr. Karl Z. Morgan, Director, Health Physics Division, Oak Ridge National Laboratory; Dr. Victor P. Bond, Brookhaven National Laboratory; Dr. George V. LeRoy, Associate Dean, School of Medicine, The University of Chicago; Dr. C. L. Comar, Chief, Biomedical Research, Oak Ridge Institute of Nuclear Studies; Dr. Stanton H. Cohn, Head, Internal Toxicity Branch, U. S. Naval Radiological Defense Laboratory, San Francisco 24, California; Mr. Charles Sandhaus, University of California, Berkeley, California.

DOE ARCHIVES

[REDACTED]

TABLE ONE

RELATIVE DOSES TO GASTROINTESTINAL TRACT
FROM INGESTION OF FALLOUT MATERIAL

	<u>Time After Detonation</u>		<u>Limiting</u>
	<u>1st Hour</u>	<u>1st Day</u>	<u>Case *</u>
Lower Large Intestine	1.0	1.0	1.0
Upper Large Intestine	1.3	0.71	0.49
Small Intestine	0.26	0.054	0.03
Stomach	0.86	0.063	0.03

*Based on assumption that there is no significant decrease in activity during time of passage through gastrointestinal tract. After a week following a detonation the decrease in activity between the mid-point of time in lower large intestine is within 20% of this condition.

DOE ARCHIVES

[REDACTED]

[REDACTED]

TABLE TWO

APPROXIMATE FISSION PRODUCT ACTIVITIES

(MICROCURIES PER MILLILITER OF GRAM $\times 10^2$)

TO PRODUCE ONE RAD DOSE TO LOWER LARGE INTESTINE*

<u>Duration of Ingestion</u> (Days)	<u>Start of Intake</u> (Days after detonation)							
	1 (1st Hour)	2 (24th Hour)	3	4	5	10	15	20
1	35	2.5	1.9	1.7	1.4	1.1	1.1	1.0
2	24	1.7	1.1	0.89	0.81	0.62	0.57	0.53
3	15	1.3	0.82	0.65	0.56	0.41	0.40	0.37
4	13	1.0	0.65	0.53	0.46	0.33	0.30	0.29
5	12	0.9	0.57	0.44	0.39	0.28	0.25	0.22
10	9.2	0.64	0.40	0.29	0.25	0.17	0.14	0.13
15	7.8	0.53	0.33	0.26	0.21	0.13	0.11	0.097
20	7.5	0.49	0.29	0.21	0.18	0.11	0.089	0.079

* a. Activities computed at start of intake period.

DOE ARCHIVES

b. Based on intake of 2200 milliliters or grams of water and food per day for adults.

~~TOP SECRET~~

TABLE THREE

CONCENTRATIONS IN WATER ON ISLANDS
IN THE PACIFIC AND ESTIMATED GAMMA
DOSE RATES AT D + 1, THREE FEET ABOVE
GROUND

Rongelap Island
(3.5 roentgens per hour)

<u>Date</u>	<u>Location</u>	<u>Gross Fission Product</u> <u>Activity (d/m/ml)</u>
D + 2	Cistern - Rongelap Islands	~ 50,000 - 75,000
D + 34	" "	~ 5,500
D + 34	Openwell "	~ 2,000
D + 300	Cistern "	~ 3
D + 330	" "	~ 4
D + 600	" "	~ 5.5
D + 600	Openwell "	~ 0.5
D + 600	Cistern (With collapsed roof)	~ 1.3

Kabelle Island
(19 roentgens per hour)

D + 330	Ground water	~ 48
---------	--------------	------

Eniwetok Island
(8.5 roentgens per hour)

D + 330	Cistern	~ 25
---------	---------	------

Eribuk Island
(1.3 roentgens per hour)

D + 600	Standing water from can, drum, etc.	~ 1.4
---------	-------------------------------------	-------

DOE ARCHIVES

TABLE FOUR

SOME POSSIBLE BIOLOGICAL EFFECTS FROM RADIATION DOSES

TO SPECIFIC ORGANS *

<u>Dose</u> <u>(Rads)</u>	<u>Gastrointestinal</u> <u>Tract</u>	<u>Thyroid</u>	<u>Bones</u>
10,000		Minor changes in structure	
	Permanent or serious damage -- survival threatened		Tumor production
1,000	Tumor Production		
	Immediate effects such as nausea and vomiting	Potential carcinogenic dose to thyroids of few percent of children and adolescents	Minor changes in structure
100			DOE ARCHIVES

* Lesser short term effects would be expected from the same doses distributed in time.

OF [REDACTED]

TABLE FIVE

MEAN BODY BURDEN OF RONGELAPESE

<u>Radioisotopes</u>	<u>Estimated Activity at One Day (μC)</u>
Sr ⁸⁹	1.6 - 22
Ba ¹⁴⁰	0.34 - 2.7
Rare earth group	1.2
I ¹³¹ (in thyroid)	6.4 - 11.2
Ru ¹⁰³	0.013
Ca ⁴⁵	0.019
Fissile material	0.016 (μgm)

DOE ARCHIVES

[REDACTED]

160
[Handwritten marks]