

**Cesium-137 and Total Potassium in Pandanus collected  
at Rongelap Island during March, 1959.**

		Cs <sup>137</sup> d/m/g dried	Cs <sup>137</sup> μmc/g wet	K mg/g dried	Cesium units μmcCs <sup>137</sup> gram K
Fruit (edible portion)	#1	424	38	5.2	37000
	#2	223	20	27.2	3700
	#3	99	8.6	9.7	4600
	#4	74	6.3	11.5	2900
	#5	266	24.	7.1	17000
	Ave.		217	19	12.2
Fruit (inedible portion)	#1	284	25	7.3	17000
	#2	79	6.8	9.2	3900
	#3	126	11	13.6	4200
	#4	65	5.9	14.0	2100
	#5	174	15.	8.4	9400
	Ave.		146	13	10.6
Leaves	#1	80	12	0.90	40000
	#2	67	10	6.2	4900
	#3	46	6.8	4.2	5000
	#4	35	5.5	10.5	1500
	#5	10	1.5	1.8	2840
	Ave.		48	10	4.7
Leaves, seedlings		343	52	3.7	42000

**Cesium-137 and Total Potassium in Pandanus collected  
at Rongelap Island during March, 1959.**

		Cs <sup>137</sup> d/m/g dried	Cs <sup>137</sup> μmc/g dried	K mg/g dried	Cesium units μmc Cs <sup>137</sup> gram K
Old bark	#2	22	10	0.80	13000
New bark	#2	30	14	5.8	2400
	#3	42	19	4.3	4500
	#4	38	17	4.3	4100
Green stem	#2	83	38	6.0	6200
	#4	59	27	10.2	2600