

4/10

Sr<sup>90</sup> in Land Plants  
Collected at Eniaetokj - March 1958  
(Exclusive of pandanus)

Sample	Sample No.	Sr <sup>90</sup> d/u/g/dry	Sr <sup>90</sup> d/u/g/wet	mg Ca gm dry sample	$\frac{\mu\text{c Sr}^{90}}{\text{gm Ca}}$	Limit of detectability
Tacca Corm peelings	R-122B	21.4 $\pm$ 2.4	3.9 $\pm$ 0.4	5.261	1900	
Cucurbita fruit Meat	R-108	35.1 $\pm$ 1.7	5.5 $\pm$ 0.27	5.943	2700	
Fimbristylis Leaves	R-116	62.3 $\pm$ 2.6	27.8 $\pm$ 1.1			
Morinda Leaves	R-118	138.4 $\pm$ 6.0	21.0 $\pm$ 0.41			
" "	R-105	99.6 $\pm$ 4.1	18.3 $\pm$ 0.76			
Cucureita, fruit meat (squash)	R-110	4.8 $\pm$ 1	0.7 $\pm$ 0.1	5.94	360	
Cucureita, seed pulp	R-109	not detectable	6.505	0	0.3 d/m/g wet	
Tacca, corn peeled	R-122c	not detectable	not detectable	1.845	0.1 d/g/wet	
Cassytha, whole plant	R-120	3.4 $\pm$ 0.5	0.7 $\pm$ 0.1	19.5	79	