

Sr<sup>90</sup> in Land Plants  
 Collected at Kabelle[- March 1958  
 (Exclusive of pandanus)

Sample	Sample No.	Sr <sup>90</sup> d/m/g/dry	Sr <sup>90</sup> d/m/g/wet	mg Ca gm dry sample	$\frac{\mu\text{C Sr}^{90}}{\text{gm Ca}}$	Limit of detectability
Guettarda Leaves	R-78	185.6 ± 6.3	36.7 ± 1.2			
" "	R-48	208.8 ± 8.1	56.0 ± 2.2			
" "	R-64	93.7 ± 4.0	26.0 ± 1.1			
Morinda Leaves	R-45	468.0 ± 18.6	88.6 ± 3.5			
" "	R-76	318.0 ± 12.4	55.3 ± 2.2			
" Fruit	R-46	54.0 ± 2.4	8.7 ± 0.4			
Lepturus Leaves	R-47	167.0 ± 6.4	----	17.016	4500	
" "	R-92	319.3 ± 13.1	154.1 ± 6.3	11.660	12000	
Fimbristylis Leaves	R-58	49.4 ± 2.3				
Tacca Tuber, Central Part	R-93C	54.7 ± 2.6	22.6 ± 1.1	4.212	5900	
Letturus, leaves	R-55	336 ± 1.9	-	15.62	9700	-
Tacca, corn unpeeled	R-93c	2.6 ± 0.4	1.1 ± 0.2	4.212	280	-