

FEDERAL GOVERNMENT

Office Memorandum • UNITED STATES GOVERNMENT

TO : Richard Hirsch, US Navy

FROM : Morse Salisbury, Director
Division of Information

SUBJECT: IODINE 131

SYMBOL: IS:MS

R

Attached is a draft for a report to be included in the 1954 annual Report of the Atomic Energy Commission, with reference to recovery of Iodine-131 from the atmosphere following the Pacific test. This was discussed at the meeting of the Committee on the Production of Radioisotopes and Related Problems.

We suggest that this be discussed in the context of the annual report. Any comments should be made by July 24.

The. 131, 1954, 1954

REPRODUCTION BY THE DWIGHT D. EISENHOWER LIBRARY

OCB File 55

FEDERAL GOVERNMENT

Richard Hirsch
Director
Division of Information
Atomic Energy Commission
Washington, D.C.
Aug 1954

Fall-out in the United States

Following nuclear detonations, radioactive debris is distributed by normal air currents over large areas, and with sufficiently sensitive instruments may be found to encircle the globe. Small amounts were deposited widely over the United States during the Pacific tests and in some areas resulted in transitory rises of the normal background radiation levels.

Transportation of the radioactive materials to the United States took only several days. Thus some of the shorter half-life radioisotopes, such as Iodine 131 (8-day half life), were still present in the fall-out. Although the amounts were biologically insignificant, it was possible, by special techniques, to demonstrate radioiodine in the thyroid glands and in the urine of grazing animals. Extremely minute quantities of Iodine 131 were also detectable in the urine of some humans for a short time.

The radioactive isotopes to be found normally in the body are potassium 40, carbon 14 and radium 226. The radiopotassium and radiocarbon are distributed throughout the tissues while the radium is almost entirely located in the skeleton. In addition to this internal irradiation, man is subjected to cosmic rays from without and to the gamma rays from radium in the soil. To this natural exposure, the radiation from bomb products is added. The point of interest in terms of health lies not in the mere presence of radioisotopes, but in the amounts and more specifically in the quantity of radiation doses delivered by these radioisotopes. The levels of activity from fall-out, outside the area surrounding the Pacific Proving Grounds, have been far less than any required to produce detectable injury either from the radioisotopes within the body or from external radiation.