

of two more adolescents, two young adults, and one man approaching middle age were sufficiently different from normal to cause Drs. Conant and Reil to describe them as questionable.

To summarize, there are now eleven of eighty-two Marshallese with definite abnormalities of their thyroid tissues and five more are considered questionable.

BEST COPY AVAILABLE

HDBruner :pwg
10/10/65

CLDunham
10/ /65

10/ /65

10/ /65

July 12, 1965

MEMORANDUM FOR: CHAIRMAN SEABORG
COMMISSIONER PALFREY
COMMISSIONER RALCY
COMMISSIONER TAPE

THROUGH GENERAL MANAGER

SUBJECT: THYROID DISEASE IN RONGELAP ISLANDERS

During the week of July 5, 1965, the three Rongelapese who were brought to this country for study at Brookhaven National Laboratory were operated on at the Lahey Clinic, Boston. You will recall that thyroid tumors were noted at the 1965 medical examination of those exposed to fallout in March 1954 from the Bravo Test.

The two teen-age boys were found to have each a solitary benign tumor of the thyroid. These nodules were removed surgically. The middle-aged woman was found to have a carcinoma of the thyroid gland with some metastases to local structures. A subtotal thyroidectomy was carried out. In her case, the prognosis is fair. She will be given 30 millicuries of Iodine-131 therapeutically to destroy persisting thyroid tissue.

It is interesting that the dose of radioiodine to the thyroid gland in the case of the woman is estimated at about 160 rads, or approximately one-tenth that to the glands of the youths. Presently under consideration is the administration of desiccated thyroid tablets prophylactically to all the Rongelapese who were at Rongelap or Ailinginae at the time of fallout.

BEST COPY AVAILABLE

Charles L. Dunham, M.D.
Director, Division of
Biology and Medicine

cc: Secretariat (2)
CL
ASM/ED
DMA

BM DIRECTOR ASM/ED

GM

DUNHAM:RAE

7/12/65

D/K Navy Files