

US DOE ARCHIVES
326 US ATOMIC ENERGY
COMMISSION
BY AUTHORITY OF DOE 5650.2, III-12 (ADMIN)

BY Joseph A. Sawyer DATE 3/26/81
BY Wilbur G. Sawyer DATE 3-26-81
Collection OGM Files
Box 326-78-3 Box 2
MRA-7-3 Analysis of
Soil From Tests 1954

NON-CONF

John C. Bugher, M. D., Director
Division of Biology and Medicine

May 4, 1954

410210

Gordon M. Dunning, Health Physicist
Biophysics Branch, Division of Biology and Medicine

DATA ON SOIL SAMPLES COLLECTED ON THE ISLANDS OF THE PACIFIC FOLLOWING THE FIRST DETONATION AND ALSO THE DOSE RATE READINGS AT THE SAME LOCALITIES.

BEST COPY AVAILABLE

Attached (Annex I) are data on soil samples collected on the islands of the Pacific following the first detonation and also the dose rate readings at the same localities. One method of evaluating such data is to try to establish relationships between different units if possible. One useful relationship would be the conversion of disintegrations per minute per gram of soil to milliroentgens per hour of gamma radiation at a three-foot height or vice versa.

Larson's work with soils around HPG during the spring 1953 tests indicated the following relationships: 10 $\mu\text{c}/\text{sq. ft.}$ beta counts of soil (after absorption and geometry corrections) \rightarrow 1 mr/hr gamma at 3 feet. He found that essentially all of the activity was in the first one inch of top soil.

In collecting soil on the Pacific Islands good care was taken to collect one square foot of surface (in fact, templates were made for this purpose). It was impossible, however, to scoop up the soil to a uniform depth so the rule followed was to collect to one inch or greater. If the fallout activity in the Pacific Islands also was contained in the first one inch, the additional soil below this contributed mass but little activity to the sample. By taking the disintegrations per minute per gram (after a thorough mixing) and multiplying by the total number of grams for each sample one should arrive at the activity per square foot.

The plot of beta disintegrations per minute per gram of soil versus mr/hr of gamma at 3 feet is shown on the attached graph. The correlation is not too good. For references, several curves are arbitrarily drawn on the graph. The data strongly suggest that less than 10 $\mu\text{c}/\text{sq. ft.}$ is equivalent to 1 mr/hr . For lower levels of activity the data are more of the order of 2 $\mu\text{c}/\text{sq. ft.}$ \rightarrow 1 mr/hr and for higher levels of activity less than 1 $\mu\text{c}/\text{sq. ft.}$ \rightarrow 1 mr/hr . If one is justified in trying to fit a curve to such widely scattered data, then 1 $\mu\text{c}/\text{sq. ft.}$ \rightarrow 1 mr/hr is a very rough approximation. Another possibility but not probability is a more complex curve shown by the solid black line.

OFFICE	BMBP			
SURNAME	Dunning: Mack	<u>WMD</u>		LABORATORY RESEARCH & APPL 7-3
DATE	5/4/54			

Form AEC-318

U. S. GOVERNMENT PRINTING OFFICE 16-62761-2

US DOE ARCHIVES

Dr. John C. Dugher

-2-

May 4, 1954

Based on calculations made in Annex II it would appear that about $10 \mu\text{c}/\text{ft}^2 \rightarrow 1 \text{ mr/hr}$ gamma at 3 feet. If one accepts the figure of 2 beta emissions for each gamma photon (Effects of Atomic Weapons) then $20 \mu\text{c}/\text{ft}^2$ (beta) $\rightarrow 1 \text{ mr/hr}$ gamma at 3 feet.

Knowing the difficulties of collecting, handling, packaging, shipping and counting the samples, it is probable that some of the data are not entirely valid. The information shown in this memo suggests that further carefully controlled studies must be made before a more firm conclusion may be reached.

Attachments 3

- Annex I
- Annex II
- Graph (w/cy 1A only)

DISTRIBUTION: copy 1A - Dr. Dugher
 2A - Dr. Dunning
 3A - Bioph. Br. Reading File
 4, 5, 6A - 30M Files

cy 5A destroyed 6/8/54

OFFICE ▶					
SURNAME ▶					
DATE ▶					

ANNEX IComparison of Soil Activity to Rose-rate Readings

<u>Location</u>	<u>Date of Collection</u>	<u>No. Grams</u>	<u>c/m/gm</u> (same date)	<u>Total Act.</u> ($\mu\text{c}/\text{ft}^2$)	<u>mr/hr</u> (survey meters)
Iikiep	6 March	2,400	23,000	2.5×10^1	~3
Jeno	"	1,060	13,000	6.2	~3
Ailuk	"	2,160	23,000	2.2×10^1	3 - 15
Kojuit	7 March	1,360	30,000	1.8×10^1	3 - 10
Orned	5 March	1,325	15,000	9.0	3.5
Erikub	5 March	1,720	4,300	3.3	1.5
Kaven	6 March	1,335	5,500	3.3	1.8
Potho	6 March	1,490	2,400	1.6	~0.8
Falap	7 March	965	950	0.4	0.5
Bongelap (Northern)	8 March	703	290,000	9.2×10^2	440.0
(Central)	"	815	1,600,000	5.9×10^2	250.0
(1 mile N. Village)	"	1,630	100,000	7.6×10^1	340.0
(South Eastern)	"	1,040	140,000	6.6×10^1	220.0
Friirippu	"	810	9,000,000	3.2×10^3	2,200.0
Eniwetok	"	2,010	780,000	7.1×10^2	900.0
Kabelle	"	1,470	4,500,000	3.0×10^3	2,000.0
Utirik	9 March	1,140	1,100,000	6.9×10^2	40.0
Bikar	"	1,030	85,000	4.1×10^1	140.0
Eniwetak	10 March	1,050	195,000	8.8×10^1	280.0
Sifo	"	1,060	14,000	6.7	100.0

The following information was obtained from the
 data collected during the survey in the manner
 described in the report.

May 11, 1954

ANNEX IICalculations of Dose Rate at Three Feet Above A Plane Surface

For point source (0.3 - 3.0 Kev range)

$$\text{Dose rate (r/hr)} = \frac{6CE}{d^2}$$

where: C = activity (curies)
E = energy in Kev
d = distance in feet

$$\text{Dose rate} = \frac{6FA \int_{R_1}^{R_2} \frac{2\pi x dx}{h^2 + x^2}}$$

where: A = activity/unit area
h = height above surface (feet)
x = distance in feet

$$\begin{aligned} \text{Dose rate} &= 6FA \int_{R_1}^{R_2} \frac{2\pi x dx}{h^2 + x^2} \\ &= 6\pi FA \ln \left[\frac{h^2 + R_2^2}{h^2 + R_1^2} \right] \end{aligned}$$

The mean free path for 0.7 Kev in air is about 360 feet so that essentially all of the dose will be contributed from a surface 1,000 feet in radius.

$$\begin{aligned} \text{Let } A &= 10 \times 10^6 \text{ curies/ft}^2 \\ E &= 0.7 \text{ Kev} \end{aligned}$$

$$\begin{aligned} \text{Dose rate} &= 6\pi(0.7)(10 \times 10^6) \ln \left[\frac{9 + 10^6}{9} \right] \\ &\approx 2.2 \text{ mr/hr} \end{aligned}$$

Since this formula assumes no absorption and also a uniform plane surface, an estimate is that

$$10 \mu\text{c/ft}^2 \longrightarrow 1 \text{ r/hr.}$$

