

QUALITATIVE DISTRIBUTION OF RADIONUCLIDES AT RONGELAP ATOLL

E. E. HELD

Laboratory of Radiation Biology, University of Washington,
Seattle, Washington

In March, 1958, a radioecological study of Rongelap Atoll was instituted at the request of the Division of Biology and Medicine, U. S. Atomic Energy Commission (AEC). This report will be concerned with generalizations regarding the distribution of radionuclides at the atoll in the fall of 1959, some five years after contamination with radioactive fallout.

Rongelap Atoll was accidentally contaminated on March 1, 1954, with radioactive fallout from a thermonuclear device detonated at Bikini Atoll some 80 miles to the west. Gamma radiation dose rates at Rongelap on D + 1 (detonation + one day) ranged from 3.5 roentgens per hour at the southern islets of the atoll to 35 roentgens per hour at the northern islets (Dunning, 1957). Eighty-two natives residing on Rongelap Island, in the south, were evacuated and did not return until June, 1957. At that time the returning population approached 300 in number but since appears to have stabilized at 230.

Several radiological and biological surveys, primarily of a monitoring nature, were conducted from the time of the first contamination until 1958 (Dunning, 1957). During this time the gamma radiation dose rates over land areas declined at approximately the rate predicted for mixed fission products by Miller and Leeb (1958). Slight rises in gamma dose rate were observed in 1956 and 1958, resulting from tests conducted during these years. However, the total contribution of radionuclides from these subsequent fallouts amounted to a fraction of one per cent of the amount from the 1954 fallout.

Rongelap Atoll is located in the Marshall Islands, in the Central Pacific Ocean, at about 11° North. It is a typical atoll with a lagoon area of 388 square miles and about 180-foot average depth. The emergent land area is about three square miles and is made up of 61 small islets ranging in size from a fraction of an acre to the largest island, Rongelap, which is about four miles long and one-half mile across at its widest point.

There is one small islet on the western reef and the remainder are strung along the northern, eastern, and southern reefs. The islets on the northern reef are not as well developed as those to the east and south. The waters of the lagoon are essentially isothermal (Robinson, 1954). The circulation, generated by the northeast trade winds, is from east to west at the surface with a returning bottom current (Von Arx, 1954). The estimated time for renewal of water in the lagoon is about 30 days.

The parent material of both soils and the lagoon bottom is primarily calcium carbonate originating mainly from coralline algae, corals and foraminifera. There is also some accumulation of humic drift in the soils.

The natives of the area are Micronesians. Their agriculture is limited to a variety of products and the only significant export is copra. About half of the food consumed at the present time is imported. Fish and other marine organisms are eaten but these sources are not exploited as much as they could be. The Rongelapese are almost en-

tirely dependent on cisterns as a source of water. Rainfall in this area is comparatively low and the islets small, so that there is not a well-developed fresh water lens. There is, however, some potable water in wells at Rongelap and Eniwetok Islets.

The native style wattle and palm frond buildings have been replaced by plywood and aluminum structures built to Rongelapese specifications by the AEC. Sanitation habits have been altered by the advent of pit toilets.

The terrestrial fauna is limited in variety. The only mammal present is the small field rat, *Rattus exulans*. The most common birds are the fairy tern, *Gygis alba*, and the noddy terns, *Anous stolidus* and *A. tenuirostris*, which nest in large numbers on some of the uninhabited islets. The reptiles are represented by skinks, geckos, and a blind snake. Land crabs are common, the most spectacular being *Birgus latro*, the coconut or robber crab. Insects are few, both in number of species and individuals. The most severe pest appears to be the beetle, *Brontispa* sp., which attacks the coconut palm.

In contrast to the land areas there is a tremendous proliferation of both numbers and variety of organisms on the reefs and in the lagoon. For example, there are over 700 species of fish. Plankton, however, is extremely sparse and as a consequence the water is so clear that green algae are found growing at depths of 180 feet.

Since the question of the effects of radiation on the organisms inevitably arises, it might be well to consider it briefly before going on to the main subject. There is no doubt that the levels of radiation were of sufficient intensity to affect living organisms. However, under actual field conditions and without benefit of study before the addition of radiation as an ecological factor, it is difficult to do more than speculate concerning the cause of the specific anomalies observed. Fosberg (1959) has accurately described the poor condition of the plants at the northern islets of Rongelap Atoll and has suggested that the primary cause of this condition is radiation. In our opinion, however, other factors, particularly edaphic factors, have probably been more important than radiation. The fact that the nitrogen content of the soils of the northern islets is lower than that of the rest of the atoll is at least circumstantial evidence that for some time there have been differences between these areas with respect to plant growth. Stone et al. (1957) have concluded from studies of *Drosophila* populations at Bikini, Eniwetok, Rongelap, and uncontaminated atolls that while there is evidence of genetic changes caused by radiation other factors mask the radiation effects. In short, it is not likely that such questions will be resolved without controlled experimentation with the species involved, under varying conditions, and with an eye toward the possibility of synergistic effects.

Approximately five years after fallout the long-lived fission products cesium-137 and strontium-90 are the principal radionuclides found in the land organisms, while the neutron-induced radionuclides zinc-65, cobalt-60 and manganese-54 are found primarily in the marine organisms.

Still detectable in the soil are manganese-54, iron-55, cobalt-57, cobalt-60, zinc-65, strontium-90, zirconium-95, ruthenium-106, antimony-125, cesium-137, cerium-144, and europium-155, which remain concentrated in the upper one to two inches. Where higher levels of radionuclides have been present these nuclides have been reported in a wide variety of organisms. It is likely that most of these radionuclides are actually present in most if

BEST COPY AVAILABLE

167

not all organisms at Rongelap but that the levels at which they occur are extremely low and so escape detection.

Passing from the soil to the soil solution, the term being used here to mean leachates collected in the field from lysimeters, strontium-90, cesium-137, and antimony-125 are the principal nuclides found, although Ru¹⁰⁶-Rh¹⁰⁶, cerium-144, and europium-155 are also detectable (Cole et al., 1961). Here differences exist with respect to soil type in that the leachates from immature soil, consisting almost exclusively of parent material, contained only antimony-125 and strontium-90. There is to us no evident explanation for this difference.

The ground water probably contains these nuclides since their movement has been detected in leachates to depths of 30 inches, but the levels are so low in ground water that special techniques would have to be developed to detect them.

The land plants contain principally cesium-137 and strontium-90. Manganese-54 and zinc-65 have been found in plants from the more heavily contaminated islets but are present in relatively insignificant amounts. In general, cesium-137 accounts for 90 per cent or more of the radioactivity in the land plants and strontium-90 for the remainder. This is unlike the situation usually found on continental soils and is a consequence of the low potassium content of Rongelap soil. Amendments of potassium to Rongelap soil reduce the uptake of cesium-137 by plants (Walker et al., 1961), and affect the distribution of cesium-137 within the plant. There are, of course, differences between plant species and plant parts with respect to the relative amounts of cesium-137 and strontium-90. For example, copra contains very little strontium-90 as compared with Pandanus fruit, and the basal leaves of various plants contain more strontium-90 relative to cesium-137 than do the terminal leaves. This variation is related to differences in mobility between cesium and potassium, and strontium and calcium.

The rats contain cesium-137 and strontium-90, reflecting the radionuclides present in the plants on which they feed. The coconut crab and the land hermit crab (*Coenobita perlatus*) contain the same nuclides but concentrate strontium-90, as has been reported for *Coenobita* from Eniwetok Atoll (Held, 1960).

The occurrence of radionuclides in man at Rongelap has been summarized by Cohn et al. (1960). In 1958 these nuclides were cesium-137 and strontium-90 coming from the food plants, and zinc-65 coming to man from marine products.

The birds, which feed almost exclusively on marine organisms, contain primarily zinc-65 and occasionally small amounts of manganese-54 and cobalt-60. Strontium-90 is also found in small amounts in bird bone and may reflect direct uptake from the ingestion of soil, although there is no direct evidence that this occurs.

Radionuclides in fish are limited to manganese-54, cobalt-60, and zinc-65, the latter being predominant. On a dry-weight basis for a sample of goatfish (*Mulloidichthys samoensis*) testes have the highest levels, the liver, gastrointestinal tract, and eyes are lower by about an order of magnitude, and the muscle and bone lower by still another order of magnitude. If the total amount of radioactivity by tissue is considered, then bone is the principal depository of zinc-65 (Joyner, 1961, personal communication). The sources of zinc-65 for fish are open to question. In some instances in-

vertebrates containing zinc-65 are known to be consumed by fish found to contain zinc-65, but, in general, no definite sources of zinc-65 are known to exist five years after fallout. It is possible that there is concentration of undetectable levels from the sea water or algae. The possibility that most of the zinc-65 radioactivity in fish is residual appears to be ruled out by the fact that young fish contain relatively high levels.

The marine invertebrates taken as a whole contain a wider spectrum of radionuclides than do the fish. These are manganese-54, cobalt-57, 60, zinc-65, strontium-90, cerium-144, and probably europium-155. The corals contain cobalt-60 and are the only invertebrates in which strontium-90 has been consistently detected. From limited data available thus far it appears that these nuclides were deposited in the skeletal material soon after fallout and have remained localized in portions of the coral colony actively growing at that time. The clams contain mostly zinc-65, cobalt-57 and cobalt-60. Weiss and Shipman (1957) originally reported the concentration of cobalt-60 in the kidney of Tridacnid clams collected at Rongelap in 1956. Animals such as the sea cucumber (*Holothuria*), *Stichopus* and spider snail (*Lambis*, *Strombus*), which ingest large amounts of bottom sediments, contain ruthenium-106, cerium-144, and probably europium-155.

Of several species of algae sampled in 1959 the only radionuclides detected were ruthenium-106, cerium-144 and europium-155. In general, the levels of radioactivity in the algae are lower than in the fish or invertebrates.

The plankton contain manganese-54, cobalt-57, 60, zinc-65, zirconium-95, ruthenium-106, and cerium-144, but all in minute amounts. In 1959 plankton samples collected by pumping a total of two and a half million gallons of water were pooled for gamma-ray spectrum analysis and were found to contain only enough of these nuclides for qualitative analysis without resorting to chemical separations. Further analysis has been deferred until other studies with the individual samples can be completed.

The lagoon sediments contain strontium-90, ruthenium-106, cerium-144, and europium-155. The radioactivity is associated mainly with the fines and is concentrated in the top two to four inches, dropping off rapidly with depth.

Radionuclides other than naturally occurring potassium-40 were not detected in sea water although larger samples and more sensitive techniques undoubtedly would have revealed their presence.

In sum, on land the present distribution of long-lived fission products, strontium-90 and cesium-137, can be expected to remain very much as it is now. The levels of radioactivity will be reduced primarily by physical decay of the radionuclides so long as other factors such as changed agricultural practices or a catastrophic storm do not occur. In the lagoon, the levels of radioactivity will decline more rapidly than on land because of the presence of shorter-lived radionuclides, with the exception of strontium-90. The latter does not enter the marine food web to any significant extent and may remain as a label useful in evaluating the long-term effects of physical forces in the lagoon.

SUMMARY

The qualitative distribution of radionuclides at Rongelap Atoll is determined approximately five

years after contamination by fallout from a thermonuclear device indicates distinct differences between the terrestrial and marine environments. The levels of radioactivity are low, the concentrations being less than the maximum permissible concentration for radionuclides in food or drinking water of man. Of the wide spectrum of radionuclides concentrated in the surface layers of the soil, strontium-90, antimony-125, and cesium-137 are the principal nuclides entering into the soil solution. The principal nuclides in the land plants and plant-eating animals such as coconut crabs and the indigenous rats are cesium-137 and, to a lesser degree, strontium-90. Bottom sediments contain mainly strontium-90 and europium-155. The radionuclides in the lagoon water have not been detected but are probably present in minute amounts. Planktonic organisms contain traces of manganese-54, cobalt-57,60, zinc-65, zirconium-95, ruthenium-106 and cerium-144. The principal nuclide found in the marine algae is cerium-144. In the marine invertebrates cobalt-60 and zinc-65 occur most commonly. Corals and coralline algae contain some strontium-90, while the fish and sea birds are found to contain mostly zinc-65. The presence of zinc-65, cesium-137, and strontium-90 in the body of the natives reflects a diet of both marine and terrestrial origin.

ACKNOWLEDGMENTS

This work was performed under contract number AT(45-1)540 between the U. S. Atomic Energy Commission and the University of Washington.

REFERENCES

- Cohn, S.H., J.S. Robertson, and R.A. Conard. 1960. Radioisotopes and environmental circumstances: The internal radioactive contamination of a Pacific island community exposed to local fallout. In R.S. Caldecott and L.A. Snyder (eds.), Radioisotopes in the Biosphere, Univ. of Minnesota Printing Dept., Minneapolis. pp. 306-330.
- Cole, D.W., S.P. Gessel, and E.E. Held. 1961. Use of the tension lysimeter in coral atoll and glacial till soils. Soil Sci. **25**: 321-325.
- Dunning, G.M. (ed.). 1957. Radioactive contamination of certain areas in the Pacific Ocean from nuclear tests. U. S. Atomic Energy Commission, U. S. Government Printing Office, Washington, D.C. 60 pp.
- Fosberg, F.R. 1959. Long-term effects of radioactive fallout on plants. Atoll Research Bulletin No. 61. National Academy of Sciences-National Research Council, Washington, D.C. 11 pp.
- . 1959. Plants and fallout. Nature **183**(4673): 1448.
- Held, E.E. 1960. Land crabs and fission products at Eniwetok Atoll. Pacific Sci. **14**(1): 18-27.
- Miller, C.F., and P. Loeb. 1958. Ionization rate and photon pulse rate of fission products from slow neutron fission of U²³⁵. U. S. Naval Radiological Defense Laboratory, San Francisco, report USNRDL-TR-247. 91 pp.
- Robinson, M.K. 1954. Sea temperature in the Marshall islands area. U. S. Geological Survey professional paper 260-D. pp. 281-291.
- Stone, W.S., M.R. Wheeler, W.P. Spencer, F.D. Wilson, J.T. Neuenschwander, T.G. Gregg, R.L. Seecof, and C.L. Ward, 1957. Genetic studies of irradiated natural populations of Drosophila. In Studies in the genetics of Drosophila. Univ. of Texas, Publ. 5721. pp. 260-316.
- Von Arx, W.S. 1954. Circulation systems of Bikini and Rongelap lagoons. U. S. Geological Survey professional paper 260-B. pp. 265-273.
- Walker, R.B., E.E. Held, and S. P. Gessel. 1961. Radiocesium in plants grown on Rongelap Atoll soils. In Recent Advances in Botany, Proc. of the 9th Intern. Bot. Congr., Univ. of Toronto Press, Toronto, Canada. pp. 1363-1367.
- Weiss, H.V., and W.L. Shipman. 1957. Biological concentration by killer clams of cobalt-60 from radioactive fallout. Science **125**(3250): 695.