18 March 1954

HEADQUARTERS JOINT TASK FORCE SEVEN APO 187 (HOW) c/o POSTMASTER

SAN FRANCISCO, CALIFORNIA

1-3/729.3

SUBJECT: Radiological Surveys of Several Marshall Island Atolls

326 US ATOMIC ENERGY RG

COMMISSION

410645

TO:

Distribution

Location

Collection

1. Attached herewith for your information and retention are copies of radiological surveys made on certain Marshall Island Atolls. The surveys were conducted as a result of contamination deposited on the affected atolls by BRAVO Shot, Operation CASTLE, fired from a reef approximately one and one half nautical miles southwest of Namu, Bikini Atoll. BRAVO Shot time was 1845 Zebra, 28 February 1954.

2. Water and soil samples were shipped to the Health and Safety Laboratory, New York Operations Office, Atomic Energy Commission (Attention: Mr. Merrill Eisenbud) for analysis.

FOR THE COMMANDER:

BEST COPY AVAILABLE

DISTRIBUTION:	(M. ammi)
CTG 7.1	- Copy 1-30 E. McGINLEY
CTG 7.2	- Copy 31 Brigadier General, U.S. Army
CTG 7.3	- Copy 32 Chief of Staff
CTG 7.4	- Copy 33
crg 7.5	- Copy 34
CINCPAC	- Copy 35
CINCPACFLT	- Copy 36
HICOMTERPACIS	- Copy 37
COMNAVSTAKWAJ	- Copy 38 /
DM:/AEC	- Copy 39
DBM/AEC	- Copy 40
Ch AFSMP	- Copy 41
CG FldComd(DWET)	- Copy 42 CLASSIFICATION CANCELLED
C/S USA, Exchat	- Copy 43 BY AUTHORITY OF DOE OG
LASL H_Div	- Copy 44 /x(uan 10/17/89)
HISL, NYOO (c/o Mgr Opns)	- Copy 45-46 , EVEWED BY QUAL DATE 125 43
USS RENSHAW (DDE-499)	- Copy 49
USS PHILIP (DDE-498)	- Сору 48
USS NICHOL:S (DDE-449)	- Copy 49
3 Incles	
וייייינוג כ	

1. Report on Soil and Water Sampling Mission by Maj R. D. Crea

2. Report on Soil and Water Sampling Mission by Dr. T. N. White, LISL

3. Rad. Survey of Downwind Atolla Contaminated by Brake by Dr. Herbert Scoville

213-54E

CA CONTRACTOR

HEADQUARTERS
JOINT TASK FORCE SEVEN
APO 187 (HOW) c/o POSTMASTER
SAN FRANCISCO, CALIFORNIA

CCTT

8 March 1954

SUBJECT: Report on Soil and Water Sampling Mission

TO:

Commander
Joint Task Force SEVEN
APO 187 (HOW)
c/o Postmaster
San Francisco, California

1. In compliance with your oral instructions, the undersigned visited LIKIE and AHUK Atolls, JEMO Island and MEJIT Island in the Eastern Marshalls between the period 5-8 March 1954 for the purpose of collecting soil and water samples and measuring level of gamma radiation present at those places in connection with BRAVO. The mission, consisting of the undersigned and a Marshallese interpreter, Lan Lakapun, embarked on the USS RENSHAW (DDE499) at Kwajalein, visited the four sites and returned to Bikini, where the remainder of the trip to Eniwetok was performed by PBM. There follows a detailed discussion of the findings at each location:

a. LIKTEP ATOLL. The samples were taken on Likiep Island, which had the largest native population. Access to the lagoon was gained through South Pass. Poor light at the end of the day and numerous coral heads necessitated anchoring about 4 miles from Likiep Island. Trip in was made by whaleboat the following morning. A water sample was taken from a large cistern fed from the roof of the Catholic rectory, and earth samples were taken from random spots about the islandable were unsheltered by trees or other growth at approximately 0800 M 6, March 1954. Radiation readings were taken with a MX-5 instrument between 0800 M and 0900 is and showed a maximum of 3 milliroentgens per hour. No variations from this reading were noted on clothing or bare feet of individuals. According to accounts received by Bishop Feeney, S.J., the population was greatly excited by the light and blast wave, the latter which reportedly arrived about 30 minutes subsequent to the light flare. According to Bishop Feeney, church attendance was greatly stimulated on the day of the test.

b. JEMO Island. This location was reached at 1100 M, 6 March 1954. It consists of a small heavily wooded island, surrounded by a line coral reef with heavy surf on three sides. There being no place for landing a whaleboat, personnel and equipment were transferred from the whaleboat to the reef by a one man rubber raft. The undersigned transferred himself by swimming. The island proved to be uninhabited, and reportedly is a sea turtle preserve. Turtle hunters erected several houses, a rain barrel of which provided a water sample. Earth samples were gathered at random from open areas, including one of beach sand above the high tide mark. The party was led straight across the island and back to the landing area via the beach, in order to verify its uninhabited state. Samples were

PAGE 151 52-213-54-1E

SECRET


RECTRICTED BAT

COLPT
SUBJECT: Report on Soil and Water Sampling Mission

collected at approximately 1200 M, 6 March 1954. Instrument readings with the MX-showed a maximum of 3 mr/hr, however this was not considered reliable, since a higher scale showed a lower reading.

- c. AILUK ATOLL. The ship reached this stell at approximately 1600 M, 6 March 1954, and slowly moved to an anchorage off Ailuk Island, the most heavily populated. The lagoon has not been swept, and numerous coral heads and pinnacles provided considerable hazard to ship movement. The landing party moved ashore by whaleboat without difficulty, and again obtained water samples from the most promisent cistern and soil samples from random unsheltered spots. Readings with the Mishowed approximately 3 mr/hr (off the 2 mr scale). An AN/PDR-27E showed a high reading of 7 mr/hr, however, on a different scale a reading of 12 or 15 mr/hr was obtained. The MX-5 reading is probably nearest correct. No significant variation were detected on bare feet or clothing of individuals. Samples and readings were taken at approximately 1700 M, 6 March 1954.
- d. <u>MEJIT Island</u>. This single coral island is also surrounded by a reof, as is JEMO, but landing was possible with a whaleboat, due to an area protected fi the surf. The island was found to be heavily populated in view of its sizd, the total number of people being 327, according to the island magistrate. Soil and water samples were taken as in the previously described manner, at approximately 1300 M, 7 March 1954. Readings with the HX-5 showed maximum of approximately 3 mr/hr (off the 2 scale, but approximately 1.5 on the 20 scale); the maximum reading the PDR 27 E was 10 mr/hr. The true figure was probably somowhere between the two.
- 2. <u>CONCLUSIONS</u>. Low level (less than 10 mr/hr) radiation measurements with field instruments of the type used are highly unsatisfactory. One MX-5 and thre: AN/PDR 27 E instruments all showed widely variant readings on different scales, and varied among each other when exposed to the same radiation. An AN/PDR T1-B proved completely useless not holding to zero even after an hours warm-up, and also showing widely variant readings on different scales.
- 3. RECOMENDATIONS. Landing parties in islands such as JEMO and MEJIT shoul be provided with a rubber 6-man or 8-man pneumatic boat, to provide greater safety to personnel and equipment. This will permit landing directly on live coral reefamith less danger of the boat being stove in. Ships assigned to such missions should draw such equipment prior to departure.
- 4. The successful accomplishment of the mission was greatly facilitated by the interest and enthusiasm of the Commanding Officer of the USS RENSHAW, CDR L. Elford, USN, and his officers and men. Their material contributions were necessar to the mission, however, the many valuable suggestions and assistance in solutions of problems proved invaluable.

/s/ R. D. Crea R. D. CREA MAJ, USA



MENORANDUM FOR: CJTF SEVEN



10 March 1954

SUBJECT: Report on Soil and Water Sampling Mission

- 1. In compliance with your oral instructions, the undersigned visited Wotj Erikub, Maloelap, Wotho, Majuro Atolls in the Marshall Islands 5 through 7 March 1954 for the purpose of obtaining earth and drinking water samples, and of measuring gamma ray dose rates, and also checked the radiological condition of the S.S ROQUE on its arrival at Majuro 7 March 1954.
- 2. The first four atolls were visited by Marshallese interpreter Takushi and the writer by means of an UF-1 amphibious aircraft. Majuro was reached by C-47. Erikub might have been emitted since it was not inhabited, being property of the Wotje tribe which goes there only occasionally to gather copra. (This was unknown until after the visit.)
- 3. At each atoll, only the principal inhabited island was visited. At each visited island an effort was made to compose a representative soil average by collecting into a single container several samples, each approximately one square foot of area and one inch depth. Water samples were collected from the principal sources currently in use. The gamma dose rates are averages for the inhabited areas.
- 4. With regard to certain minor discrepancies between the survey methods up by Major R. D. Crea and the writer; it was originally planned to perform the surjointly, and when it became advisable to separate and survey different atells, no time remained for discussion of details of techniques.
- 5. Gamma-ray dose rates on Wotje and on Erikub are each the average of MX-and N/PDR-39 average readings which agreed reasonably well. The MX-5 was rende inoperative when the rubber life raft was swamped by surf on the first attempt to launch from the beach at Erikub. Following the Wotho survey, the PDR-39 developed temperature-dependent reading of 0.4 2 mr/hr, so that later readings in this range are of very dubious reliability.
- 6. The following tabulation summarizes the atoll survey. S is Soil, W is Water Sample:

ATOLL	ISLAND	DATE	TIME	SAMPLE NO	MR/HR & SAMPLING
WOTJE	ORMED	5 Mar	1600	S 5	3.5 mr/hr, 1-beach, 3-mid-vill
				W6	age, 1-back village. ½ well plus ½ catch basin.
ERIKUB	ERIKUB	5 Mar	1715	S6	1.5 mr/hr. 1-mid-village, 1 on path to boach. No inhabitants, no water supply found.









ATOLL	ISLAND	DATE	TIME	SAMPLE NO	MR/HR & SAMPLING
MALOELAP	KVAEN	6 Mar	1130	37	1.8 mr/hr, 2-village, 2-path to beach.
				W12 W13	Well water. From catch basin.
OHTOW	WOTHO	6 Mar	1615	S8	0.8 mr/hr, 1 by well; 2-mid- village.
		٠.		W9	Well water (no rain in catch basin for 2 mo.)
MAJURO	ULIGA	7 lar	1200	59	0.5 mr/hr, 4 from near Admin Bldg.
				W10	Tap water.

- 7. Pacific Micronesian Line S.S. "ROQUE", Master: Lawrence Blane, home port Guam, left Ebeye 0840 M on 1 March, entered channel to Utirik Lagoon about 1200 M on 2 March, and anchored in Lagoon at 1524 M on 2 March; docked at Majuro (Uliga Is.) 1630M on 7 March. Readings (mr/hr) after docking: 2-3 inside main deck struture, 10 on open deck, 5-8 in sleeping quarters on upper deck, 10-30 on rope and canvas. Prior radiation levels cannot be estimated because of rain squalls and u certainty about when decks last washed. Laster was advised to have decks washed down as soon as convenient. He was told that the activity would not hurt hayone, but that it was undesirable to have it around longer than necessary.
- 8. RECOMMENDATIONS: Future visits to Erikub and Maloclap should not be attempted by UF-1 except under conditions of greater urgency. The writer's prior experience in such operations is very limited, but from his own observations plus the remarks made by those better qualified to judge, it appears that a fair amous of risk is involved.
- 9. Especially notable was the very cooperative attitude of the Navy personat Kwajaloin and the Marshall District Administrative Officials at Majuro in supporting this mission.

l Incl:
Marshall Islands Atoll
Samples collected by T. N.
White, 5-7 March 1954

/s/ T. N. White DR. T. N. WHITE Health Division LASL








MARSHALL ISLANDS ATOLL SAMPLES COLLECTED BY T. N. WHITE, 5-7 MARCH 1954


Earth samples were collected as follows:

At each island visited several samples were dug and put into the same one-gallon "ice-cream carton". Each sample (i.e. each digging) approximated one square foot to a depth of one inch. The number and locations of the samples were selected to represent, as well as could be judged, an average of the areas used by the inhabitants, after the samples were mixed in the carton. Areas that wer unusually shaded or unshaded by trees were avoided. The large "pebbles" in the composite represent coral gravel from "main street" through the village.

Water samples were selected according to the principal source in current us

Inclosure 1





HEADQUARTERS TASK UNIT 13
Task Group 7.1
APO 187 (HOW) P.O. Box 8
c/o Postmastor
San Francisco, California

TU-13-54-375

12 Merch 1954

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

1. Acknowledgement

The members of the survey team wish to express their appreciation to Captain, officers and members of the crow of the USS NICHOLAS (DDE 449) for the assistance and cooperation in conducting the survey herein reported. Captain Elliot turned over all possible facilities of his ship in order to assist in the survey. LT Frink, the Executive Officer, organized all the operations of the beat parties, and it was only through his personal direction and participation that it was possible to carry out the small beat surveys under extremely difficenditions. Since most of the lagoon waters were not navigable by a DDE, it we necessary to make long beat trips in high seas and land on tricky coral recess. That it was possible to make, without mishap, a detailed survey of five widely separated atells in the course of three days with only two beats was largely detection is efforts.

2. Introduction

The BRAVO Shot contaminated a number of atolls in generally eastward direction from Bikini to such an extent that it became necessary to evacuate the native populations from Rongelap, Alinginae and Utirik Atolls and the military personnel on Rongerik Atoll. Following this evacuation CJTF SEVEN organized to subject detailed radiological survey of the atolls to the eastward of Bikini (Ref. CJTF SEVEN Enimetok 060400Z). The data from this survey were required for the following purposes:

- a. The evaluation of the radiation effects on evacuees.
- b. The estimation of the clapsed time before reoccupancy.
- c. The estimation of the residual radiation effects of large yields surface detonations.

In connection with this survey, teams from various Task Groups and Mr Wilds, Trust Territory Representative, returned to the atolls to secure the evacuated habitations, service military equipment, and obtain documentary phot graphy.

3. Operational Schedule

8 Harch - 0800 Survey team rendezvous aboard USS NICHOLAS (DDE 449) in Rongelap Legoon.



SECRET

236

213-54-3E

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

- 8 March 1000 1800 Two parties in small boats surveyed living arcs on Rongelap Island and castorn half of Rongelap Atoll
- 9 March 0700 1130 Two parties in small boats proceeded from the I which was stationed outside Utirik Atoll and surveyed Utirik and Aon Islands, the main islands of the Atoll
- 9 March 1500 1700 One party in a small boat landed on the outer reef of Bikar Island and surveyed the island, the onl large island of Bikar Atoll.
- 10 March 0700 1100 Two parties in small boats proceeded from the D which was stationed outside Rongerik Atoll and survey Eniwetak Island (where the Task Force's Units had been stationed) and the other important islands of the Ato
- 10 March 1430 1980 Two parties in small boats proceeded from the D which was stationed outside Alinginae Atoll and survey the inhabited islands of the Atoll.
- 11 March 0700 1400 One party in a small boat survoyed the northwoorn islands of Rongelap Atoll and one party rechecks the living areas on Rongelap Island and established a reference location for future decay-measurements.
- 12 March 0800 Survey toam arrived Eniwetok Atoll via DDE.

4. The following personnel from test projects in TG 7.1, TU 13, served as members of the survey team:

Horbort Scaville, Jr.	TU-13 Staff
Richard Rast	Project 2.1
Richard Soulc	Project 2.5a
Walmer Strape	Project 6.4

The USS NICHOL'S (DDE 449) supplied boat crews under the direction of LT Cliffor Frink, Executive Officer, for surveys.

5. Instrumentation

Radiac set M/PDR-39 was selected as the instrument to be used in the conduct of the survey. Five (5) each of AN/PDR-39 were calibrated with an 80 Curic Co⁶⁰ source twenty-four hours before departure. The calibration yielded zero variation between instruments - any scale. Upon cross checking three of these instruments, (a point of actual survey) in a radiation field of 0.320 r/mit was found that all three instruments gave the same reading.

These survey meters were subject to prolonged use under adverse conditions of dampness (to the point of sea water splashing over them), salt deposit and continual rough handling. With one exception, all instruments operated efficiently for the duration of the operation. On the final day it was found

that one survey meter could not be properly zero adjusted. The four remaining AN/PDR-39, still operated efficiently and seemed to be in good working order.

- One (1) each Bockman MX-5, and one (1) each /N/PDR-27A was brought alon for any low intensity checks nocessary. Two (2) each calibrated AN/PDR-TLB, were on hand to serve as spares in the event of operational failure with the /N/PDR-3: None of these instruments were required.
- 6. The average and maximum gamma dose rates measured on the various island of each atoll are plotted in Figures 1 through 5. All measurements were made at waist height unless otherwise indicated. The maximum readings do not include measurements made with the instrument next to a contaminated surface.

Detailed surveys were made of all the inhabited localities. Typical readings are given in Tables 1 and 2 for the native village of Rengelap Island, and the TG 7.4 camp on Eniwetak Island. In general, the villages and the camps appeared to have slightly lower average dose rates than the remainder of the island. This can perhaps be ascribed to different geometry of the contamination and to slightly greater penetration into the losse gravel in the native villages. The dose rates inside the native huts appeared to be almost the same as the dose rate outside. The dose rate in the middle of the military barracks, tents, and shacks was 1/3 to 1/2 that outside. This reduction is probably largely a geome' effect. The dose rate fell off rapidly on the beach below the high tide mark. There was no evidence of rain washing off the contaminated material. The feilage on the windward sides of the islands appeared to be slightly above average contamination.

TABLE 1

TYPICAL READINGS IN RONGELAP VILLAGE - 8 MARCH

Location	Dose Rate (mr/hr)
Rongelap Island (average) Center of village Near central cistern Near southern cistern Near northern cistern	375 280 300 220 350

TABLE 2

TYPICAL READINGS IN CAMP ON ENTWETCH IS. - 10 MARCH

Location	Outside Dose Rate (<u>cr/hr)</u>	Rate (mr/hr)
Eniwetak Island (average) Mess hall Tent, edge of main camp Latrine Sleeping quarters Dispensary	280 220 270 - 260 260 220	110 175 160 90
	U22.13.1	238

•	- ·	
Radio Station	290	160
Weather Station (N end of island	1) 280	110
Proj 6.6. Station (S end of islan	nd) 240	<u> </u>

In order to estimate the rate of decay between 8 and 11 March, the following radiation measurements were taken on three days on Rongelap Island:

	8 March	11 March
Central living area (village) Southern most cistern Roof of cistern (Southern most) Ground (contact) cistern area	280 mr/hr 220 mr/hr 240 mr/hr 220 mr/hr	170 mr/hr 145 mr/hr 140 mr/hr 110 mr/hr

An area was selected 30 yards inland from the Rongelap cemetery as a measing point for future decay measurements. This area is outlined with 2X4s place on pails. The waist height reading was 210 mr/hr at 1000 hours, 11 March 1954.

7. Sample collections

Water samples were collected from the water supplies of all inhabited areas. About two quarts of water were transferred to a polyethlene bottle at each site. These will be turned over to the New York Operations Office, AEC analysis.

Soil samples were collected at all inhabited areas and also at several uninhabited islands. In collecting the soil samples a one foot by one foot square was marked on the ground and soil to about one inch of depth was removed from the square and transferred to a cardboard container. The primary samples will be turned over to the New York Operation Office, ARC, for analysis, and some smaller samples will be analyzed by Program 2 of TU 13.

Listed in Table 3 are the samples taken with the dose rate measured at waist height at the location where they were taken.

TABLE 3 - SOIL

Sample No.	<u>Atoll</u>	Island	Date	Mr/Hr
1*	Rongelap	Rongelap (North end)	8 Mar	440
<u>2</u> 3	Rongelap	Rongelap (Center of village)	8 Mar	280
3	Rongelap	Rongelap (1 mile north of village)	8 Mar	340
4	Rongelap	Rongelap (near South cistern of village)	8 Har	2 20
5 *	Rongelap	Eriirippu	8 Mar	2200
6 *	Rongelap	Eniaetok	8 Mar	900
7 *	Rongelap	Kabelle	8 Mar	2000
8*	Utirik	Utirik	9 Mar	40
9	Bikar	Bikar	9 Mar	160
10	Rongerik	Eniwetak	10 Mar	280
11*	Ailinginae	Sifo	10 Mar	100

*Small additional sample taken for analysis by Program 2 of TU 13.



7

TABLE 3 - WATER

Sample No.	Atoll	Island	Date	Ur/Hr
_1-	Rongelap	Rongelap (central cistern)	8 Mar	300
_2	Rongelap	Rongelap (North part of village)	8 Mar	350
3	Rongelap	Rongelap (Northernmost cistern)	8 Mar	400
4	Rongelap	Rongelap (Southernmost cistern)	8 Mar	220
5	Utirik	Utirik (cistern near church)	9 Mar	40
6	Utirik	Utirik (cistern at south of village)	9 Mar	40
7	Rongerik	Eniwetak (Distillation water)	10 Mar	240

In addition to the above, a sample of foilage was taken at the wind ward side of Bikar Island. The radiation field was 180 mr/hr on 9 March 195; at this point.

8. Conclusions and Recommendations

- a. The radiological survey proved that a large yield surface detontion can produce extremely serious radiological contamination over a distance more than 120 miles downwind and important contamination about 250 miles downwind.
- b. The center of the contamination pattern from the BRAVO Shot lies somewhat north of Rongelap and Rongerik Atolls and probably not far from a libetween Bikini and Bikar.
- c. Although the fall—out was serious on Rongelap Island located at the extreme southeast tip of the atoll, the contamination was about ten times greater at the north side of the atoll, twenty miles away.
- d. The contamination decreased by a factor of about eight over the downwind distance of 50 miles between Rongelap and Rongerik.
- e. Standard military field housing provides a significant degree of protection to personnel inside.
- f. The AN/PDR-39 proved to be a very satisfactory instrument for field survey work under rigorous environmental conditions.
- g. A single DDE with two (2) whale boats is not a completely satisfactory method of conducting a broad radiological survey of the type just completed. Future surveys should consider using vessels capable of entering more of the atolls and of handling a helicopter and several small boats.

6 Incls:

1. Rad. Survey Rongelap

2. Rad. Survey Utirik

3. Rad. Survey Bikar


4. Rad. Survey Rongerik


5. Rad. Survey Ailinginae

6. Summary of Rad. Survey

/s/ Herbert Scoville
DR. HERRERT SCOVILLE
Technical Director
AFSVP


ncehill ild onn







242

فخذ







BIKARA	XX	AILUK 20 20 20 20 20 20 20 20 20 20 20 20 20	
SUMMARY OF RADIOLOGICAL SURVEY NOTE ALL VALUES ARE ROBBY, EXTRA- POLATED BY MEANS OF 717 BCCAY LAW THESE VALUES DIO NOT ACTUALLY XINST SINCE TIMES OF AMEN'AL OF COMTAWINANT WERE CONSIDERABLY LATER.	TAKA	AIL.	
	AGERIK		5.0.6
	A STANDARY WAS THE WAS		
3.09	RONGELAP RONGELAP 10 PURCHEALER 10 PURCHEALE	WOTHO B.	
	CHARLE FOR	N. C.	,