

482

410717

546.211

# On the Distribution of Radioactivity in the Sea around Bikini Atoll in June, 1954.

by

Y. Miyake, Y. Sugiura and K. Kameda  
Meteorological Research Institute

Received January 6, 1955

## Abstract

By the radioactivity in the air and sea water caused by the US hydrogen bomb tests in the area of Bikini-Eniwetok Atolls, 1954, the radioactive contamination to fishes and fishing boats were brought about. The observation of the actual state of the radioactivity around Bikini-Eniwetok Atolls was carried out aboard the survey boat "Shunkotsu-maru". The maximum radioactivity in sea water was 7025 cpm/l at 570 km west of Bikini, and the influence of the radioactivity was recognized as far as about 2000 km from Bikini Atoll. Some oceanographic discussions on the horizontal as well as vertical distributions of the radioactivity in the sea are given.

## 1. Introduction

Leaving Tokyo Harbour on 15th May, 1954, the observation were carried out until 4th July aboard the survey boat the "Shunkotsu-maru" belonging to the Ministry of Agriculture and Forestry in the sea area near Bikini Atoll where H-bomb tests had been done by U.S. Atomic Energy Commission, to make clear the state of the radioactive contamination to living matter, sea water, the air etc. The purpose of this "Japanese Bikini Expedition" was to obtain a counter-plan against the serious damage fell on the Japanese pelagic fisheries due to the radioactivity carelessly scattered in this sea area since the unfortunate affair of the fishing boat the "Fuku-ryu-maru", the "Lucky Dragon" on 1st March, 1954.

In this expedition the present authors took charge of making a plan and observations on the radioactive contamination of sea water and the air. Two of the authors (SUGIURA and KAMEDA) participated in the expedition as observers. The fairly strong radioactivity was detected in wide area around Bikini Atoll, though U.S. Atomic Energy Commission had declared that the presence of the radioactivity in the area was negligible [1]. In this paper the authors intend to report the results of observations and some oceanographic discussions regarding the distributions of the radioactivity near Bikini Atoll.

## 2. The method of measurement of radioactivity in sea water

WNRC

It is difficult to measure the radioactivity in sea water directly by simple evaporation method owing to its high salinity, therefore, the carrier method has been adopted. After adding 2 gr of solid ammonium chloride, 1 ml of the aqueous solution of iron alum (5.3 gr/l) and 1 ml of barium chloride solution (17.8 gr/l) to 1 litre of sea water, water is heated to 60~70°C stirring well. A few drops of an alcoholic solution of phenol-phthalein (1%) are added as an indicator, then ammonium

BEST COPY AVAILABLE 27

79-0121 Box 3  
#25 REDWING  
FALLOUT RESULTS

hydroxide solution 1:1 is carefully dropped in with a pipet until a faint pink colour appears. After two minutes boiling, the precipitate is settled on standing for a couple of hours at room temperature. The precipitate is filtered under suction on a round filter paper (dia. 30mm) laying above the glass filter (Fig. 1). The precipitate remained on a filter paper is dried in an air bath or a desiccator holding in a small brass case (Fig. 2) to avoid the folding of the surface. Measurements of radioactivity are done by putting the precipitate kept in a brass case under a mica window of G-M counter. The G-M counter used aboard the "Shunkototsu-maru" is Radiation Counter, Model 32 manufactured by Science Research Institute Ltd., Tokyo. The thickness of mica window is  $3.7 \text{ mg cm}^{-2}$  and the distance between the mica window and the surface of the precipitate is 1.5 cm. After the measurement has been done, the precipitate is covered by a polyethylene film to protect the surface. It was confirmed that at least 70-80% of the total activity in sea water could be transferred to the precipitate by preliminary tests using the fission materials fell aboard the fishing boat the "Fukuryu-maru". Therefore, the actual activities might be 25-40% higher than observed values. To obtain the counting efficiency the aqueous solution containing  $\text{Ce } 144$  of which activity had been referred to the




Fig. 1. The apparatus for filtration.




Fig. 2. A small brass case to avoid the folding of the surface of a filter paper.

standard RaE was treated chemically as mentioned above, and its radioactivity was measured under the same geometrical conditions. Thus, the efficiency of the measurement was found to be 7.7%, that is, 1000 counts per minute is equivalent to about  $5.9 \text{ m}\mu\text{C}$ . The maximum activity in sea water obtained aboard by this method was  $7025 \text{ cpm/l}$  which was found at the depth of 75 m, at 450 km west of Bikini Atoll on 21st June.

The second was  $6050 \text{ cpm/l}$  found at the depth of 20 m at 150 km west of the atoll on 12th June. It was confirmed that most parts of the radioactive materials were dispersed in water as true solutions, ionic or eu colloidal, since more than

99% of the radioactivity passed through a filter paper (pore size  $0.5 \mu$ , No. 4 Filter Paper, Tōyō Filter Paper Manufacturing Co., Tokyo).

It is rather surprising that the activity in sea water near Bikini and Eniwetok Atolls was often stronger than, or at least comparable to, that of White Oak Lake receiving waste effluent from nuclear reactors of Oak Ridge National Laboratory USA. The latter was 2710~5190 cpm per litre on an average [2].

3. The distribution of radioactivity in sea water around Bikini Atoll

a. Horizontal distribution

Fig. 3 shows the horizontal distribution of the activity in the surface water around Bikini Atoll. There must have been some changes in the activity due to the time variation of the sea condition, the natural decay and the loss of the radioactivity during the course of observation. However, since the causes of changes were much complicated, and the correction were almost impossible, the distribution was drawn by using directly the observational data without giving any correction.



Fig. 3 The horizontal distribution of the radioactivity of the surface water. Number expresses the radioactivity in cpm/l.

As shown in Fig. 3 the radioactive water extended on the WNW direction leaving near Bikini Atoll while a branch spread out to WSW. The cause of this branching is not clear. But, as explained below, it seems that there was a weak discontinuous boundary along the line passing Bikini and Eniwetok Atolls. The water  $B_1$  (Fig. 4) on the northern side of this line had on an average the




Fig. 4. The course lines of observation, the water systems, and the current speed and its direction.

temperature, 27.1°C, the chlorinity, 19.23‰ and  $\sigma_t$ , 22.51 at 50 m depth. On the other hand, the water system  $B_2$  on the southern side of the line had T, 27.0°C, Cl, 19.04‰ and  $\sigma_t$ , 22.29. Thus, it seems that there was a slight but definite jump in the properties between two water systems, which might be one of the causes of the branching.

The radioactive water flowing in the main direction of the North Equatorial Current extended sooner toward the west than in the southern part where the flow rate was smaller because the latter was near the boundary between the North Equatorial Current and the Equatorial Counter Current and there was also the oppression due to the north-eastward inclination of the Equatorial Counter Current. Thus, on the WNW direction the activity of 100 cpm  $I$  reached as far as 2000 km from Bikini Atoll, while on the WSW direction the extension was only 1000 km.

It is to be noted that the radioactivity was also detected at the position 700~800 km from Bikini on the ENE direction. This is the end part of the radioactivity distribution turning to the east at about 350 km north of Bikini Atoll (Fig. 4). To clarify the cause of this turning on the backward direction of the North Equatorial Current may be one of the important problems.

Though it is not clear at present, whether it was the extension of the easterly current along the northern boundary of the North Equatorial Current which was found recently between 180°E~140°W (3), or it was only the local eddy, several discussions will be done below on this point.

When we investigate the horizontal distribution of water systems near Bikini Atoll by using the T-S diagrams of waters at the depth of 50 m below the surface, we can find out that there are at least four different types of waters (Fig. 4). The first one A is highly saline water (chlorinity, 19.3‰, water temperature, 27.5°C,  $\sigma_t$ , 22.77) covering the area northern than 15°N.

In the southern part of the atoll there is the second type of water C which belongs to the Equatorial Counter Current with smaller chlorinity (Cl, 18.9‰) and higher temperature (T, 28.0°C). The third one B belongs to the North Equatorial Current representing the intermediate property between A and C. In the southern part of the water C, there extends the water D of the South Equatorial Current with higher chlorinity (Cl, 19.6‰) and higher temperature (T, 28.0°C).

The water B may be classified into two sub-groups  $B_1$  and  $B_2$  as mentioned above. As shown in Fig. 4 the east part of B was comparatively broad and it became narrower on approaching to the west. It was probably due to that at the time the boundary between waters B and C somewhat inclined to the north. Therefore, the North Equatorial Current was pressed by the Equatorial Counter Current and the backward current flowing toward the north-east direction of Bikini Atoll would have been resulted.

WNRC

The observation of the current direction obtained *in situ* also suggests the presence of counter or eddy current in the same area. It is, however, not sure whether this current would form the origin of the general easterly current as mentioned above.

On the other hand, a remarkable loop of the isochlor lines shown in the surface distribution of the chlorinity (Fig. 5) in the northern area of Bikini also indicates the presence of eddy flow.

As shown in Figs. 6 and 7, in the vertical distributions of phosphate and the dissolved oxygen along 15°N, there was the subsidence in the area near 15°N.

165°E which was probably due to the anticyclonic eddy.

b. Vertical distributions of the radioactivity in sea water

Figs. 8, 9 and 10 show the distributions of radioactivity in sea water on the vertical section perpendicular to the North Equatorial Current along the lines respectively at 150 km, 570 km and 1000 km west of Bikini. The activity at 150 km west of Bikini was already the branching of the activity in the direction of N-S. The dotted lines in the figures represent respectively the latitude of the sites of Bikini and Eniwetok Atolls. The strong activity is separated on both sides of these dotted lines and the distance of two maxima is about 100 km. The depth which shows the activity of 100 cpm/l is at about 80 m below the surface in the northern and southern parts and in the middle part it sinks down to about 120 m. It is noteworthy that the activity decreases rapidly down to only 28 cpm/l at 200 m depth. This corresponds to the depth of thermocline which was about 150 m in this area.

The estimated amount of flow of radioactivity was about  $1 \times 10^5$  curie/hour passing through this section. On the vertical section at 570 km west of Bikini the distance of branching became broader and also the depth of 100 cpm/l activity was 200 m below the sea surface on an average. The flow of radioactivity was about  $0.6 \times 10^5$  curie/hour. The distance between the former and the latter section was about 400 km and assuming that the speed of current was about 0.7 knots




Fig. 7. The surface distribution of the chlorinity in ‰.




Fig. 8. The vertical distribution of the saturation rate of the dissolved oxygen ( $\sigma_x$ ) along 15°N.

WNRC




Fig. 7. The vertical distribution of phosphate P in  $\mu\text{g atoms/l.}$

Fig. 8. The vertical distribution of the radioactivity in sea water in the section perpendicular to the North Equatorial Current along the line at 150 km west of Bikini Atoll.





Fig. 9. The vertical distribution of the radioactivity in sea water in the section perpendicular to the North Equatorial Current along the line at 570 km west of Bikini Atoll.




Fig. 10. The vertical distribution of the radioactivity of sea water in the section perpendicular to the North Equatorial Current along the line at 1300 km west of Bikini Atoll.

WNRG

on the direct perpendicular to the line of the two sections would have taken about 2 weeks to flow between them. Thus, it is reasonable to think that although both sections would have been about the same when the rate of seaway during two weeks was taken into consideration.

#### 4. The relation between the radioactivity in sea water and the radioactive fallout

The fishing boat "Shunkotsu-maru" was situated at the position 150 km east of Bikini Atoll on the early morning on 1st March when her crews recognized small dust particles falling on the deck. The density of dusts covered the surface of the deck was not clearly known, however, it may be supposed that it is about 0.5 gr/m<sup>2</sup> or so. On the other hand, the radioactivity of the dust particles was regarded as about 1.4 curie/gr at the time of falling (3 hours after the detonation [4]). Assuming that the mean diameter of dust particles was 0.1 mm, the rate of falling of such particles in sea water is about 30 m/hour. Then, it takes about 20 seconds to pass the water layer of 1 cm. Here, if we assume that about 0.1% of the radioactive substances in the dusts could dissolve into sea water in one second, most part of radioactive materials would have removed to water from dusts before they reached 100 m depth below the sea surface. Thus, the radioactivity in water near the sea surface around there would have been about 10<sup>-4</sup> curie/l at the time five or six hours after detonation. Needless to say, this was the activity of sea water in the area about 150 km apart from Bikini Atoll, therefore, the activity near and in the atoll would have been much more stronger. On the other hand, it was confirmed by the records of micro-barographs and tide gauges obtained in Japan [5] that the detonations had been carried out four or five times since 1st March (1st March, 17th March, 6th Apr., 25th Apr. and 5th May) it would be no wonder the wide sea area far from Bikini Atoll had been contaminated by fission materials as strongly as it was actually observed aboard the "Shunkotsu-maru" in June, 1954.

#### 5. The property on the radioactive substances in sea water

The relation between the gross radioactivity,  $A_t$ , of the fission materials and the time after detonation may be simply expressed by the next equation,

$$A_t = A_0 e^{-\alpha t}$$

where  $c$  and  $\alpha$  are constants. From the decay curve of the radioactivity of the precipitates prepared aboard the "Shunkotsu-maru", it has been found that  $\alpha$  is confined within the comparatively narrow limits of 1.3~1.6 (mean, 1.5) assuming the date of detonation was 5th May. In normal cases,  $\alpha$  is about 1.2[6], whereas it was 1.37 for the radioactive dusts fell on the deck of the "Fuku-ryu-maru".

In sea water some of the radioactive elements contained in the original fission materials might be preferentially precipitated owing to pH of sea water, (about 8.2~8.3). Therefore, the chemical composition should be somewhat different from the original fission materials, which might be the cause of the change in the value of  $\alpha$ .

The radiochemical analysis of the fission materials in sea water have been furnished also by the present authors which will be reported in near future. In Table 1 the full data of the gross activity of sea water observed aboard the observation vessel are given.

*Acknowledgements* The authors wish to express their thanks to Dr. H. YABE, the leader of the Expedition, Messrs. G. YOSHII, H. MIYOSHI, Y. YAZAKI who carried out the radioactivity measurement aboard the ship, and also to members of laboratories of Dr. K. KIMURA and Dr. E. MINAMI at Tokyo University for their help and cooperation.

#### References

- [1] U.S. Atomic Energy Commission, 1950. Radiobiological Survey of Bikini, Eniwetok and Likiep Atolls. *Health, 2*, 187. AECU-3446.
- [2] Radiotope Uptake by Aquatic Plants and Animals Living in Waste Effluents of Oak Ridge National Laboratory. A paper presented to U.S.-Japan Radiobiological Conference held in Tokyo, 1954.
- [3] CROMWELL, T., 1953. Circulation in a Meridional Plane in the Central Equatorial Pacific. *Jour. Mar. Res.*, 12, 1-196.
- BARNETT, C.A., BORG, S.D.F. and LYMAN, J., 1948. Ocean Circulation in Marshall Island Area. *Trans. Amer. Geophys. Union*, 29, 6.
- [4] KIMURA, K. *et al.*, 1954. On the Radioactive Substances Having Precipitated on the Fishing Boat from the F-15, *Yokohama Japan Analyst*, 3, p. 335. (in Japanese).
- [5] YAMAMOTO, R., 1954. The Microbarographic Oscillations Produced by the Explosions of Hydrogen Bombs. *Bull. Inst. Chem. Res. Kyoto Univ. The Radioactive Central Meteorological Observatory, Tokyo*, 1954. A Preliminary Report of Observations on the Atmospheric and Oceanic Oscillations caused by H-Bomb Tests. (in Japanese).
- [6] HENNER, H.F. and BAILEY, N., 1951. Fission-Product Decay Rates. *Nucleonics*, 9, p. C-1.

WNRG



Date	St. No.	Latitude Longitude	Depth m	Activity cpm/l	Date	St. No.	Latitude Longitude	Depth m	Activity cpm/l	
June, 20	21-4	38°30'N 160°21'E	0	42	June, 20	13	38°03'N 159°53'E	0	16	
			12	48				26	7	
			48	56				30	9	
			72	57				42	6	
			78	57				49	5	
	90	57	49	3						
	4	38°5 160°2'5		24	36	14	14	38°24.5 159°31'	0	0
				48	48				20	0
				82	70.25				49	0
				103	162				72	9
144				23	106				0	
21-4	38°42 160°15		0	30	15	25	38°59 159°51'	0	10	
			25	46				25	10	
			49	39				49	0	
			91	46.4				73	0	
			216	0				95	tr.	
21	22	38°31 160°36	0	20.7	21	26	38°00 159°10	0	10	
			12	26.6				25	7	
			25	22				50	9	
			48	22				71	69	
			72	20				72	7	
			80	6				93	tr.	
			10	4.4				485	tr.	
			200	0				0	0	
			280	0				0	0	
			360	0				0	0	
405	0	0	0							
485	22	0	0							
715	40	0	0							
21-4	39°18.5 160°5		0	4.7	21	28	38°10 160°08.5	0	7	
			25	31.5				27	3	
			50	22				51	3	
			75	2.7				74	3	
			84	2.4				110	3	
21-4	39°05 158°57		0	18.9	21	29	38°00 160°21	0	0	
			25	20.7				25	7	
			48	13.4				50	10	
			75	3.4				74	5	
			99	0				98	0	
198	0	198	0							
22	4	39°3 158°41	0	0	22	4		25	0	
			25	35				50	10	
			49	30.4				74	5	
			73	0				98	0	
			95	0				198	0	
			175	0				525	0	
455	2									

WARG