

# 487  
HEADQUARTERS  
JOINT TAILOR SEVEN  
APO 187 (HOW) c/o POSTMASTER  
SAN FRANCISCO, CALIFORNIA

410854

J-3/729.3

18 March 1954

SUBJECT: Radiological Surveys of Several Marshall Island Atolls

TO: Distribution

342  
Location Technical Library B-2  
Collection AFWC  
Folder Castle Survey of Marshall Islands Mar. 1954-June 1954

1. Attached herewith for your information and retention are copies of radiological surveys made on certain Marshall Island Atolls. The surveys were conducted as a result of contamination deposited on the affected atolls by BRAVO Shot, Operation CASTLE, fired from a reef approximately one and one half nautical miles southwest of Namu, Bikini Atoll. BRAVO Shot time was 1845 Zebra, 28 February 1954.

2. Water and soil samples were shipped to the Health and Safety Laboratory, New York Operations Office, Atomic Energy Commission (Attention: Mr. Merrill Eisenbud) for analysis.

FOR THE COMMANDER:

BEST COPY AVAILABLE

DISTRIBUTION:

CTG 7.1 - Copy 1-30  
CTG 7.2 - Copy 31  
CTG 7.3 - Copy 32  
CTG 7.4 - Copy 33  
CTG 7.5 - Copy 34  
CINCPAC - Copy 35  
CINCPACFLT - Copy 36  
HICOMTERPACIS - Copy 37  
COMNAVSTAKWAJ - Copy 38  
DMA/AEC - Copy 39  
DEM/AEC - Copy 40  
Ch AFSWP - Copy 41  
CG FldComd(DWET) - Copy 42  
C/S USA, ExAgt - Copy 43  
LASL H Div. - Copy 44  
HASL, NYOO (c/o Mgr Opns) - Copy 45-46  
USS RENSHAW (DDE-499) - Copy 47  
USS PHILIP (DDE-498) - Copy 48  
USS NICHOLS (DDE-449) - Copy 49

*E. McGinley*  
E. MCGINLEY  
Brigadier General, U.S. Army  
Chief of Staff

3 Incls:

1. Report on Soil and Water Sampling Mission by Maj R. D. Crea
2. Report on Soil and Water Sampling Mission by Dr. F. N. White, LASL
3. Rad. Survey of Downwind Atolls Contaminated by BRAVO by Dr. Herbert Scoville

AFWL/NO

CLASSIFICATION CANCELLED  
BY AUTHORITY OF DOE/DOE\*

RECEIVED BY  
DATE  
Per DIA LTR (15CM) 7/10/85  
Carlisle 8/16/86

310

~~TOP SECRET~~  
~~CONFIDENTIAL~~  
~~SECRET~~

~~CONFIDENTIAL~~  
~~SECRET~~

~~CONFIDENTIAL~~  
~~SECRET~~

# 487  
[REDACTED]  
HEADQUARTERS  
JOINT TASK FORCE SEVEN  
APO 187 (HOW) c/o POSTMASTER  
SAN FRANCISCO, CALIFORNIA

COMPT

8 March 1954

SUBJECT: Report on Soil and Water Sampling Mission

TO: Commander  
Joint Task Force SEVEN  
APO 187 (HOW)  
c/o Postmaster  
San Francisco, California

1. In compliance with your oral instructions, the undersigned visited LIKIE and AILUK Atolls, JEMO Island and MEJIT Island in the Eastern Marshalls between the period 5-8 March 1954 for the purpose of collecting soil and water samples and measuring level of gamma radiation present at those places in connection with BRAVO. The mission, consisting of the undersigned and a Marshallese interpreter, Lan Lakapun, embarked on the USS HENSHAW (DDE-499) at Kwajalein, visited the four sites and returned to Bikini, where the remainder of the trip to Eniwetok was performed by PEM. There follows a detailed discussion of the findings at each location:

a. LIKIEP ATOLL. The samples were taken on Likiep Island, which had the largest native population. Access to the lagoon was gained through South Pass. Poor light at the end of the day and numerous coral heads necessitated anchoring about 4 miles from Likiep Island. Trip in was made by whaleboat the following morning. A water sample was taken from a large cistern fed from the roof of the Catholic rectory, and earth samples were taken from random spots about the island which were unsheltered by trees or other growth at approximately 0800 M 6, March 1954. Radiation readings were taken with a MX-5 instrument between 0800 M and 0900 M and showed a maximum of 3 milliroentgens per hour. No variations from this reading were noted on clothing or bare feet of individuals. According to accounts received by [REDACTED] the population was greatly excited by the light and blast wave, the latter which reportedly arrived about 30 minutes subsequent to the light flare. According to [REDACTED], church attendance was greatly stimulated on the day of the test.

b. JEMO Island. This location was reached at 1100 M, 6 March 1954. It consists of a small heavily wooded island, surrounded by a fine coral reef with heavy surf on three sides. There being no place for landing a whaleboat, personnel and equipment were transferred from the whaleboat to the reef by a one man rubber raft. The undersigned transferred himself by swimming. The island proved to be uninhabited, and reportedly is a sea turtle preserve. Turtle hunters erected several houses, a rain barrel of which provided a water sample. Earth samples were gathered at random from open areas, including one of beach sand above the high tide mark. The party was led straight across the island and back to the landing area via the beach, in order to verify its uninhabited state. Samples were

[REDACTED]  
AFW/AMO  
[REDACTED]  
[REDACTED]

COMPT

SUBJECT: Report on Soil and Water Sampling Mission

PRIVACY ACT MATERIAL REMOVED

collected at approximately 1200 M, 6 March 1954. Instrument readings with the MX showed a maximum of 3 mr/hr, however this was not considered reliable, since a higher scale showed a lower reading.

c. AILUK ATOLL. The ship reached this atoll at approximately 1600 M, 6 March 1954, and slowly moved to an anchorage off Ailuk Island, the most heavily populated. The lagoon has not been swept, and numerous coral heads and pinnacles provided considerable hazard to ship movement. The landing party moved ashore by whaleboat without difficulty, and again obtained water samples from the most prominent cistern and soil samples from random unsheltered spots. Readings with the MX showed approximately 3 mr/hr (off the 2 mr scale). An AN/PDR-27E showed a high reading of 7 mr/hr, however, on a different scale a reading of 12 or 15 mr/hr was obtained. The MX-5 reading is probably nearest correct. No significant variations were detected on bare feet or clothing of individuals. Samples and readings were taken at approximately 1700 M, 6 March 1954.

d. MEJIT Island. This single coral island is also surrounded by a reef as is JEMO, but landing was possible with a whaleboat, due to an area protected from the surf. The island was found to be heavily populated in view of its size, the total number of people being 327, according to the island magistrate. Soil and water samples were taken as in the previously described manner, at approximately 1300 M, 7 March 1954. Readings with the MX-5 showed maximum of approximately 3 mr/hr (off the 2 scale, but approximately 1.5 on the 20 scale); the maximum reading with a PDR 27 E was 10 mr/hr. The true figure was probably somewhere between the two.

2. CONCLUSIONS. Low level (less than 10 mr/hr) radiation measurements with field instruments of the type used are highly unsatisfactory. One MX-5 and three AN/PDR 27 E instruments all showed widely variant readings on different scales, and varied among each other when exposed to the same radiation. An AN/PDR T1-B proved completely useless not holding to zero even after an hours warm-up, and also showing widely variant readings on different scales.

3. RECOMMENDATIONS. Landing parties in islands such as JEMO and MEJIT should be provided with a rubber 6-man or 8-man pneumatic boat, to provide greater safety to personnel and equipment. This will permit landing directly on live coral reefs with less danger of the boat being stove in. Ships assigned to such missions should draw such equipment prior to departure.

4. The successful accomplishment of the mission was greatly facilitated by the interest and enthusiasm of the Commanding Officer of the USS RENSLOW, CDR USN, and his officers and men. Their material contributions were necessary to the mission, however, the many valuable suggestions and assistance in solution of problems proved invaluable.

PRIVACY ACT MATERIAL REMOVED

/s/ R. D. Crea  
R. D. CREA  
MAJ, USA

ARMED

MEMORANDUM FOR: CJTF SEVEN

10 March 1954

SUBJECT: Report on Soil and Water Sampling Mission

# 487

1. In compliance with your oral instructions, the undersigned visited Wotje, Erikub, Maloolap, Wotho, Majuro Atolls in the Marshall Islands 5 through 7 March 1954 for the purpose of obtaining earth and drinking water samples, and of measuring gamma ray dose rates, and also checked the radiological condition of the S.S. ROQUE on its arrival at Majuro 7 March 1954.

2. The first four atolls were visited by Marshallese interpreter Takushi and the writer by means of an UF-1 amphibious aircraft. Majuro was reached by C-47. Erikub might have been omitted since it was not inhabited, being property of the Wotje tribe which goes there only occasionally to gather copra. (This was unknown until after the visit.)

3. At each atoll, only the principal inhabited island was visited. At each visited island an effort was made to compose a representative soil average by collecting into a single container several samples, each approximately one square foot of area and one inch depth. Water samples were collected from the principal sources currently in use. The gamma dose rates are averages for the inhabited areas.

4. With regard to certain minor discrepancies between the survey methods used by Major [redacted] and the writer, it was originally planned to perform the survey jointly, and when it became advisable to separate and survey different atolls, no time remained for discussion of details of techniques.

5. Gamma-ray dose rates on Wotje and on Erikub are each the average of MX-5 and AN/PDR-39 average readings which agreed reasonably well. The MX-5 was rendered inoperative when the rubber life raft was swamped by surf on the first attempt to launch from the beach at Erikub. Following the Wotho survey, the PDR-39 developed a temperature-dependent reading of 0.4 - 2 mr/hr, so that later readings in this range are of very dubious reliability.

6. The following tabulation summarizes the atoll survey. S is Soil, W is Water Sample:

| <u>ATOLL</u> | <u>ISLAND</u> | <u>DATE</u> | <u>TIME</u> | <u>SAMPLE NO</u> | <u>MR/HR &amp; SAMPLING</u>                                                           |
|--------------|---------------|-------------|-------------|------------------|---------------------------------------------------------------------------------------|
| WOTJE        | ORMED         | 5 Mar       | 1600        | S5               | 3.5 mr/hr, 1-beach, 3-mid-village, 1-back village.                                    |
|              |               |             |             | W6               | 1/2 well plus 1/2 catch basin.                                                        |
| ERIKUB       | ERIKUB        | 5 Mar       | 1715        | S6               | 1.5 mr/hr. 1-mid-village, 1 on path to beach. No inhabit ants, no water supply found. |

PRIVACY ACT MATERIAL REMOVED

AFWL/NO

~~SECRET~~

| <u>ATOLL</u> | <u>ISLAND</u> | <u>DATE</u> | <u>TIME</u> | <u>SAMPLE NO</u> | <u>MR/HR &amp; SAMPLING</u>                   |
|--------------|---------------|-------------|-------------|------------------|-----------------------------------------------|
| MALOEELAP    | KA VEN        | 6 Mar       | 1130        | S7               | 1.8 mr/hr, 2-villago, 2-path to beach.        |
|              |               |             |             | W12              | Well water.                                   |
|              |               |             |             | W13              | From catch basin.                             |
|              |               |             |             |                  | PRIVACY ACT MATERIAL REMOVED                  |
| WOTH O       | WOTH O        | 6 Mar       | 1615        | S8               | 0.8 mr/hr, 1 by well; 2-mid-villago.          |
|              |               |             |             | W9               | Well water (no rain in catch basin for 2 mo.) |
| MAJURO       | ULIGA         | 7 Mar       | 1200        | S9               | 0.5 mr/hr, 4 from near Admin Bldg.            |
|              |               |             |             | W10              | Tap water.                                    |

7. Pacific Micronesian Line S.S. "ROQUE", : , home port, Guam, left Ebayo 0840 M on 1 March, entered channel to Utirik Lagoon about 1200 L on 2 March, and anchored in Lagoon at 1524 M on 2 March; docked at Majuro (Uluga Is.) 1630M on 7 March. Readings (mr/hr) after docking: 2-3 inside main dock structure, 10 on open deck, 5-8 in sleeping quarters on upper deck, 10-30 on rope and canvas. Prior radiation levels cannot be estimated because of rain squalls and uncertainty about when decks last washed. was advised to have decks washed down as soon as convenient. He was told that the activity would not hurt anyone, but that it was undesirable to have it around longer than necessary.

8. RECOMMENDATIONS: Future visits to Erikub and Maloolap should not be attempted by UF-1 except under conditions of greater urgency. The writer's prior experience in such operations is very limited, but from his own observations plus the remarks made by those better qualified to judge, it appears that a fair amount of risk is involved.

9. Especially notable was the very cooperative attitude of the Navy personnel at Kwajalein and the Marshall District Administrative Officials at Majuro in supporting this mission.

1 Incl:  
Marshall Islands Atoll  
Samples collected by T. N.  
Whito, 5-7 March 1954

/s/ T. N. Whito  
DR. T. N. WHITE  
Health Division  
LASL

PRIVACY ACT MATERIAL REMOVED

MARSHALL ISLANDS ATOLL SAMPLES COLLECTED BY T. N. WHITE, 5-7 MARCH 1954

Earth samples were collected as follows:

At each island visited several samples were dug and put into the same one-gallon "ice-cream carton". Each sample (i.e. each digging) approximated one square foot to a depth of one inch. The number and locations of the samples were selected to represent, as well as could be judged, an average of the areas used by the inhabitants, after the samples were mixed in the carton. Areas that were unusually shaded or unshaded by trees were avoided. The large "pobbles" in the composite represent coral gravel from "main street" through the village.

Water samples were selected according to the principal source in current use.

Inclosure 1

APWL/MO

# 487  
HEADQUARTERS TASK UNIT 13  
Task Group 7.1  
APO 187 (HOW) P.O. Box 8  
c/o Postmaster  
San Francisco, California

PRIVACY ACT MATERIAL REMOVED

TU-13-54-375

12 March 1954

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

1. Acknowledgement

The members of the survey team wish to express their appreciation to the Captain, officers and members of the crew of the USS NICHOLAS (DDE 449) for the assistance and cooperation in conducting the survey herein reported. Captain [redacted] turned over all possible facilities of his ship in order to assist in the survey. LT [redacted] the Executive Officer, organized all the operations of the boat parties, and it was only through his personal direction and participation that it was possible to carry out the small boat surveys under extremely difficult conditions. Since most of the lagoon waters were not navigable by a DDE, it was necessary to make long boat trips in high seas and land on tricky coral reefs. That it was possible to make, without mishap, a detailed survey of five widely separated atolls in the course of three days with only two boats was largely due to his efforts.

2. Introduction

The BRAVO Shot contaminated a number of atolls in generally eastward direction from Bikini to such an extent that it became necessary to evacuate the native populations from Rongelap, Ailinginae and Utiirik Atolls and the military personnel on Rongerik Atoll. Following this evacuation CJTF SEVEN organized the subject detailed radiological survey of the atolls to the eastward of Bikini (Ref. CJTF SEVEN Eniwotok 060400Z). The data from this survey were required for the following purposes:

- a. The evaluation of the radiation effects on evacuees.
- b. The estimation of the elapsed time before reoccupancy.
- c. The estimation of the residual radiation effects of large yields surface detonations.

In connection with this survey, teams from various Task Groups and Mr. Wilds, Trust Territory Representative, returned to the atolls to secure the evacuated habitations, service military equipment, and obtain documentary photography.

3. Operational Schedule

8 March - 0800 Survey team rendezvous aboard USS NICHOLAS (DDE 449) in Rongelap Lagoon.

[redacted]  
[redacted]

[redacted]  
[redacted]

AFWJ/NO

[redacted]

PRIVACY ACT MATERIAL REMOVED

[redacted]

[REDACTED]

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

PRIVACY ACT MATERIAL REMOVED

8 March - 1000 - 1800 Two parties in small boats surveyed living areas on Rongolap Island and eastern half of Rongolap Atoll

9 March - 0700 - 1130 Two parties in small boats proceeded from the [REDACTED] which was stationed outside Utirik Atoll and surveyed Utirik and Aon Islands, the main islands of the Atoll

9 March - 1500 - 1700 One party in a small boat landed on the outer reef of Bikar Island and surveyed the island, the on large island of Bikar Atoll.

10 March - 0700 - 1100 Two parties in small boats proceeded from the [REDACTED] which was stationed outside Rongerik Atoll and surveyed Eniwotak Island (where the Task Force's Units had been stationed) and the other important islands of the Atoll

10 March - 1430 - 1900 Two parties in small boats proceeded from the [REDACTED] which was stationed outside Alinginao Atoll and surveyed the inhabited islands of the Atoll.

11 March - 0700 - 1400 One party in a small boat surveyed the northwestern islands of Rongolap Atoll and one party rechecked the living areas on Rongolap Island and established reference location for future decay measurements.

12 March - 0800 Survey team arrived Eniwotak Atoll via DDE.

4. The following personnel from test projects in TG 7.1, TU 13, served as members of the survey team:

TU-13 Staff  
Project 2.1  
Project 2.5a  
Project 6.4

The USS NICHOLAS (DDE 449) supplied boat crews under the direction of LT [REDACTED], Executive Officer, for surveys.

#### 5. Instrumentation

Radiac set AN/PDR-39 was selected as the instrument to be used in the conduct of the survey. Five (5) each of AN/PDR-39 were calibrated with an 80 Curie Co<sup>60</sup> source twenty-four hours before departure. The calibration yielded zero variation between instruments - any scale. Upon cross checking three of these instruments, (a point of actual survey) in a radiation field of 0.320 r/hr it was found that all three instruments gave the same reading.

These survey meters were subject to prolonged use under adverse conditions of dampness (to the point of sea water splashing over them), salt deposit and continual rough handling. With one exception, all instruments operated efficiently for the duration of the operation. On the final day it was found

APRIL/NO

PRIVACY ACT MATERIAL REMOVED

[REDACTED] Cr2 [REDACTED] [REDACTED]

4

[REDACTED]

SUBJECT: Radiological Survey of Downwind Atolls Contaminated by BRAVO

that one survey meter could not be properly zero adjusted. The four remaining AN/PDR-39, still operated efficiently and seemed to be in good working order.

One (1) each Bockman MX-5, and one (1) each AN/PDR-27A was brought along for any low intensity checks necessary. Two (2) each calibrated AN/PDR-TLB, were on hand to serve as spares in the event of operational failure with the AN/PDR-39. None of these instruments were required.

6. The average and maximum gamma dose rates measured on the various islands of each atoll are plotted in Figures 1 through 5. All measurements were made at waist height unless otherwise indicated. The maximum readings do not include measurements made with the instrument next to a contaminated surface.

Detailed surveys were made of all the inhabited localities. Typical readings are given in Tables 1 and 2 for the native village of Rongelap Island, and the TG 7.4 camp on Eniwetak Island. In general, the villages and the camps appeared to have slightly lower average dose rates than the remainder of the island. This can perhaps be ascribed to different geometry of the contamination and to slightly greater penetration into the loose gravel in the native villages. The dose rates inside the native huts appeared to be almost the same as the dose rate outside. The dose rate in the middle of the military barracks, tents, and shacks was  $1/3$  to  $1/2$  that outside. This reduction is probably largely a geometrical effect. The dose rate fell off rapidly on the beach below the high tide mark. There was no evidence of rain washing off the contaminated material. The foiling on the windward sides of the islands appeared to be slightly above average contamination.

TABLE 1

TYPICAL READINGS IN RONGELAP VILLAGE - 8 MARCH

| <u>Location</u>           | <u>Dose Rate (mr/hr)</u> |
|---------------------------|--------------------------|
| Rongelap Island (average) | 375                      |
| Center of village         | 280                      |
| Near central cistern      | 300                      |
| Near southern cistern     | 220                      |
| Near northern cistern     | 350                      |

TABLE 2

TYPICAL READINGS IN CAMP ON ENIWETAK IS. - 10 MARCH

| <u>Location</u>           | <u>Outside Dose Rate (mr/hr)</u> | <u>Inside Dose Rate (mr/hr)</u> |
|---------------------------|----------------------------------|---------------------------------|
| Eniwetak Island (average) | 280                              | --                              |
| Mess hall                 | 220                              | 110                             |
| Tent, edge of main camp   | 270                              | 175                             |
| Latrine                   | 260                              | 160                             |
| Sleeping quarters         | 260                              | 90                              |
| Dispensary                | 220                              | 110                             |

[REDACTED]

|                                     |     |     |
|-------------------------------------|-----|-----|
| Radio Station                       | 290 | 160 |
| Weather Station (N end of island)   | 280 | 110 |
| Proj 6.6. Station (S end of island) | 240 | --  |

In order to estimate the rate of decay between 8 and 11 March, the following radiation measurements were taken on three days on Rongelap Island:

|                                 | <u>8 March</u> | <u>11 March</u> |
|---------------------------------|----------------|-----------------|
| Central living area (village)   | 280 mr/hr      | 170 mr/hr       |
| Southern most cistern           | 220 mr/hr      | 145 mr/hr       |
| Roof of cistern (Southern most) | 240 mr/hr      | 140 mr/hr       |
| Ground (contact) cistern area   | 220 mr/hr      | 110 mr/hr       |

An area was selected 30 yards inland from the Rongelap cemetery as a measuring point for future decay measurements. This area is outlined with 2X4s placed on pails. The waist height reading was 210 mr/hr at 1000 hours, 11 March 1954.

### 7. Sample collections

Water samples were collected from the water supplies of all inhabited areas. About two quarts of water were transferred to a polyethylene bottle at each site. These will be turned over to the New York Operations Office, AEC for analysis.

Soil samples were collected at all inhabited areas and also at several uninhabited islands. In collecting the soil samples a one foot by one foot square was marked on the ground and soil to about one inch of depth was removed from the square and transferred to a cardboard container. The primary samples will be turned over to the New York Operation Office, AEC, for analysis, and some smaller samples will be analyzed by Program 2 of TU 13.

Listed in Table 3 are the samples taken with the dose rate measured at waist height at the location where they were taken.

TABLE 3 - SOIL

| <u>Sample No.</u> | <u>Atoll</u> | <u>Island</u>                            | <u>Date</u> | <u>Mr/Hr</u> |
|-------------------|--------------|------------------------------------------|-------------|--------------|
| 1*                | Rongelap     | Rongelap (North end)                     | 8 Mar       | 440          |
| 2                 | Rongelap     | Rongelap (Center of village)             | 8 Mar       | 280          |
| 3                 | Rongelap     | Rongelap (1 mile north of village)       | 8 Mar       | 340          |
| 4                 | Rongelap     | Rongelap (near South cistern of village) | 8 Mar       | 220          |
| 5*                | Rongelap     | Eriirippu                                | 8 Mar       | 2200         |
| 6*                | Rongelap     | Eniaetok                                 | 8 Mar       | 900          |
| 7*                | Rongelap     | Kabelle                                  | 8 Mar       | 2000         |
| 8*                | Utirik       | Utirik                                   | 9 Mar       | 40           |
| 9                 | Bikar        | Bikar                                    | 9 Mar       | 160          |
| 10                | Rongerik     | Eniwetak                                 | 10 Mar      | 280          |
| 11*               | Ailinginae   | Sifo                                     | 10 Mar      | 100          |

\*Small additional sample taken for analysis by Program 2 of TU 13.

[REDACTED]  
TABLE 3 - WATER

| <u>Sample No.</u> | <u>Atoll</u> | <u>Island</u>                        | <u>Date</u> | <u>Mr/Hr</u> |
|-------------------|--------------|--------------------------------------|-------------|--------------|
| 1                 | Rongelap     | Rongelap (central cistern)           | 8 Mar       | 300          |
| 2                 | Rongelap     | Rongelap (North part of village)     | 8 Mar       | 350          |
| 3                 | Rongelap     | Rongelap (Northernmost cistern)      | 8 Mar       | 400          |
| 4                 | Rongelap     | Rongelap (Southernmost cistern)      | 8 Mar       | 230          |
| 5                 | Utirik       | Utirik (cistern near church)         | 9 Mar       | 40           |
| 6                 | Utirik       | Utirik (cistern at south of village) | 9 Mar       | 40           |
| 7                 | Rongerik     | Eniwetak (Distillation water)        | 10 Mar      | 240          |

In addition to the above, a sample of foliage was taken at the windward side of Bikar Island. The radiation field was 180 mr/hr on 9 March 1954 at this point.

#### 8. Conclusions and Recommendations

a. The radiological survey proved that a large yield surface detonation can produce extremely serious radiological contamination over a distance more than 120 miles downwind and important contamination about 250 miles downwind.

b. The center of the contamination pattern from the BRAVO Shot lies somewhat north of Rongelap and Rongerik Atolls and probably not far from a line between Bikini and Bikar.

c. Although the fall-out was serious on Rongelap Island located at the extreme southeast tip of the atoll, the contamination was about ten times greater at the north side of the atoll, twenty miles away.

d. The contamination decreased by a factor of about eight over the downwind distance of 50 miles between Rongelap and Rongerik.

e. Standard military field housing provides a significant degree of protection to personnel inside.


f. The AN/PDR-39 proved to be a very satisfactory instrument for field survey work under rigorous environmental conditions.

g. A single DDE with two (2) whale boats is not a completely satisfactory method of conducting a broad radiological survey of the type just completed. Future surveys should consider using vessels capable of entering more of the atolls and of handling a helicopter and several small boats.

#### 6 Incls:

1. Rad. Survey Rongelap
2. Rad. Survey Utirik
3. Rad. Survey Bikar
4. Rad. Survey Rongerik
5. Rad. Survey Ailinginae
6. Summary of Rad. Survey

/s/ Herbert Scoville  
DR. HERBERT SCOVILLE  
Technical Director  
AFSWP



**FIG. 1**




FIG. 2





FIG. 3



# AILINGINAE ATOLL

## RADIOLOGICAL SURVEY

10 MARCH

AVERAGE READING (MAX. READING)  
MR/HR.




FIG. 5




FIG. 6