

4 6 5

May 13, 1954

Dr. John C. Bugher, M.D., Director
Division of Biology and Medicine

Gordon M. Dunning, Health Physicist
Biophysics Branch, Division of Biology and Medicine

411534

ESTIMATED RADIATION DOSE TO THYROID OF NATIVES FROM RONGELAP

This memorandum is in reply to your request for an estimate of additional doses to the thyroid of the Rongelap natives due to the fact that tellurium, as a precursor to iodine, may be present in the gut after ingestion of fallout material. The tellurium, in turn, might disintegrate into radioactive iodine while in the gut, with subsequent deposition of the iodine in the thyroid.

There are some 17 radioactive isotopes of tellurium but only 7 of these are produced in fission. Of these, 6 are not of interest (4 have too short a half-life, 1 leads to stable iodine-127 and 1 leads to iodine-129 with a half-life of $2-4 \times 10^7$ years). The remaining radioisotope is tellurium-132, with a half-life of 77 hours leading to iodine-132 with a half-life of 2.4 hours. (Incidentally there is no tellurium precursor that is of interest here.)

Without having the original data of IASL I have accepted their estimate that there were ingested and/or inhaled the products of 5×10^{13} fissions, assumed they were all ingested, and then proceeded to calculate the dose to the thyroid from (a) I^{131} (b) each short-lived iodine isotope of interest and (c) the added dose coming from $T^{132}-I^{132}$. The calculations show that $T^{132}-I^{132}$ will produce an added dosage of about 26%.

The best estimated percentage absorption and deposition of iodine is yet to be determined. The best estimate I can turn up to date is still the 20% quoted in IBS Handbook 52. However, I will continue to search for additional information. In the meantime the table below indicates the magnitude of doses to the thyroid if one assumes 20, 50, and 100% absorption and deposition. Incidentally, it may be noted that the calculations below based on 20% (the number assumed by IASL) show estimated doses to the thyroid from I^{131} and from shorter lived iodine isotopes to be in good agreement with those estimated by IASL, i.e., 50 reps for I^{131} and 80 reps for short-lived iodine isotopes.

CLASSIFICATION CANCELED
DOE NSI DECLASSIFICATION REVIEW E.O. 12958
BY: SEAN D. GAMAGE DOE/NN-523

BEST COPY AVAILABLE

RG 386 / Job 1320
Collection ANA Operational Safety - McCann
Box /
Folder B. Rongelap. HENK FILES

OFFICE	EBBP			
SURNAME	DUNNING:MMack	<i>YMS</i>		MILITARY RESEARCH & APPL. 7-5
DATE	5-13-54			

DOSE TO THYROID (REPS)

	<u>Assuming 100% Retention</u>		<u>Assuming 50% Retention</u>		<u>Assuming 20% Retention</u>	
I ¹³¹	255	255	128	128	51	51
I ¹³²	27*	(54)**	14*	(27)**	5*	(10)**
I ¹³³	370	370	185	185	74	74
I ¹³⁵	60	60	30	30	12	12
I ¹³² (Te ¹³²)	185*	(370)**	93*	(185)**	37*	(74)**
Total	897*	(1109)**	450*	(555)**	169*	(221)**

* If assume that one-half of the I¹³² (half-life 2.4 hours) present in the gut is deposited in the thyroid.

** If assume all of the I¹³² (half-life 2.4 hours) present in the gut is deposited in thyroid.

Most probable estimate of ratio of doses to thyroid is:

$$\frac{I^{132,133,135} + I^{132}(Te^{132})}{I^{131}} \approx 2.5$$

OFFICE ▶					
SURNAME ▶					
DATE ▶					

May 13, 1954

ANNEXCalculations of Dose to Thyroid I^{131}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D / 1 there are 0.017 d/r/10,000 fissions

or 8.5×10^7 d/r/5 x 10^{13} fissions

or 33.3 μ c intake of I^{131}

or 1.4×10^{12} atoms intake of I^{131}

(Average energy 0.22 Mev)

$$\text{Dose (reps)} = \frac{(1.35 \times 10^{12}) * (0.22) (1.6 \times 10^{-6})}{(20)(93)} = 255 \text{ reps}$$

* Correction for biological decay.

 I^{132}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D / 1, I^{132} intake is 1.1×10^{11} atoms

The average mean energy of I^{132} is about 0.55 mev or 2.5 times that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$$(1.1 \times 10^{11})(2.5) = 2.75 \times 10^{11} \text{ atoms of } I^{131}$$

However, due to the short half-life of I^{132} (2.4 hrs) assume that the energy equivalent of 1.5×10^{11} atoms of I^{131} reaches the thyroid.

Thus, the ratio of doses $\frac{I^{131}}{I^{132}} = 9.0$

OFFICE ▶					
SURNAME ▶					
DATE ▶					

I^{133}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At $D \neq 1$, I^{133} intake is 1.24×10^{12} atoms

The average mean energy of I^{133} is about 0.36 or 163 times that of I^{131} .

Thus, the ratio of doses $\frac{I^{131}}{I^{133}} \approx 0.7$

 I^{135}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At $D \neq 1$, I^{135} intake is 2.36×10^{11} atoms.


The average mean energy of I^{135} is about 0.3 mev or 1.36 that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$$(2.36 \times 10^{11})(1.36) = 3.2 \times 10^{11} \text{ atoms of } I^{131} \text{ energy equivalent.}$$

Thus, ratio of doses $\frac{I^{131}}{I^{135}} \approx 4.2$

OFFICE ▶					
SURNAME ▶					
DATE ▶					



Assume: inhalation and/or ingestion of 5×10^{13} fissions.

At $D \neq 1$, Te^{132} intake 100 μc

or 1.5×10^{12} atoms

Assume: the time spent in the gut is 77 hrs.

Then, 7.3×10^{11} atoms of Te^{132} will have disintegrated into 7.3×10^{11} atoms of I^{132} .

The average mean energy of I^{132} is about 0.55 Mev or 2.5 times that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$$(7.3 \times 10^{11})(2.5) = 1.8 \times 10^{12} \text{ atoms of } \text{I}^{131}.$$

However, due to the short half-life of I^{132} (2.4 hrs), assume that only the energy equivalent of 1×10^{12} atoms of I^{131} reaches the thyroid.

$$\text{Thus, the ratio of doses } \frac{\text{I}^{131}}{\text{I}^{132}(\text{Te}^{132})} = 1.4.$$

DISTRIBUTION: copy 1A - Dr. Bugher
 2A - Fr. Dunning
 3A - Biophysics Branch
 4,5,6A --- BOM Files

By SA destroyed 6/14/54

OFFICE ▶					
SURNAME ▶					
DATE ▶					